

**Pest risk assessment for the European Community:  
plant health: a comparative approach with case studies**

The logo for Prima phacie features the word "Prima" in a green, sans-serif font, followed by "phacie" in a black, sans-serif font. The text is centered between two horizontal green bars. A faint reflection of the text is visible below the main text.

**Prima phacie**

**Pest Risk Assessment: Revised Test Method 4b<sup>1</sup>  
Without specific risk reduction options in place**

**December 2011**

**A. MacLeod<sup>1</sup>, M. Holeva<sup>2</sup>, H. M. Anderson<sup>1</sup> & J. Smith<sup>1</sup>**

<sup>1</sup> Food and Environment Research Agency, Sand Hutton, York, YO41 1LZ, UK

<sup>2</sup> Benaki Phytopathological Institute, 8 Stefanou Delta str., 14561 Kifissia Attica, Greece

---

<sup>1</sup> Method 4b describes risk elements using an ordinal scale of 5 categories (very low, low, medium, high, very high)

**Preface**

Pest risk assessment provides the scientific basis for the overall management of pest risk. It involves identifying pests and characterizing the risks associated with those pests by estimating their probability of introduction (entry, transfer and establishment) as well as the severity of the consequences to crops and the wider environment as a result of their introduction.

Risk assessments are science-based evaluations. They are neither scientific research nor are they scientific manuscripts. The risk assessment forms a link between scientific data and decision makers and should express risk in terms appropriate for decision makers.

**Note**

Risk assessors will find it useful to have a copy of International Standards for Phytosanitary Measures No. 5, the Glossary of Phytosanitary Terms (IPPC, 2007)<sup>1</sup>, ISPM No. 11, Pest risk analysis for quarantine pests, including analysis of environmental risks and living modified organisms (IPPC, 2004)<sup>2</sup> and the EFSA guidance document on a harmonized framework for pest risk assessment (EFSA, 2010)<sup>3</sup> to hand as they read this document and conduct a pest risk assessment.

---

<sup>2</sup> ISPMs Nos. 5 and 11 available at [https://www.ippc.int/index.php?id=ispms&no\\_cache=1&L=0](https://www.ippc.int/index.php?id=ispms&no_cache=1&L=0)

<sup>3</sup> EFSA Journal 2010, **8**(2),1495-1561, Available at <http://www.efsa.europa.eu/en/scdocs/doc/1495.pdf>

## CONTENTS

	<u>Page</u>
<b>Executive Summary</b>	4
<b>Stage 1: Initiation</b>	9
<b>Stage 2: Pest risk assessment (outline)</b>	14
• Pathways	15
• Potential for entry and transfer on each pathway	55
• Overall potential for entry and transfer	55
• Potential for establishment	56
• Spread	70
• Consequences of establishment and spread	75
• Potential impact	80
• Pest risk ( <i>potential for entry and transfer x potential impact</i> )	81
• Uncertainties	82
<b>Conclusion</b>	82
<b>Supporting documentation</b>	
• Annex 1: Questionnaire template distributed to EFSA Plant Health Network	

## Executive Summary

*Provide a technical summary reflecting the content of the assessment (the questions addressed, the information evaluated, and the key issues which resulted in the conclusion)*

The purpose of this pest risk assessment was to evaluate the plant health risk associated with *Acidovorax citrulli* within the framework of EFSA project CFP/EFSA/PLH/2009/01.

### **Pest biology** (see datasheet for details)

- Identity of the pest

*Acidovorax citrulli* (Schaad *et al.*, 1978) Schaad *et al.*, 2008, comb. nov.

Betaproteobacteria: Burkholderiales: Comamonadaceae


Synonym: *Acidovorax avenae* subsp. *citrulli* (Schaad *et al.*, 1978) Willems *et al.*, 1992

Schaad *et al.* (2008) proposed that the taxonomic position of the pathogen be elevated from subspecies to species level (*i.e.* *A. citrulli* instead of *A. avenae* subsp. *citrulli*). The new name was then validly published in the International Journal of Systematic and Evolutionary Microbiology in 2009 (Schaad *et al.*, 2009) However, this new nomenclature has not yet been adopted by the scientific community (Bahar & Burdman, 2010). In this assessment, the most recent name, *Acidovorax citrulli* is used.

- Life history / Disease cycle and epidemiology: (*briefly*)

*Acidovorax citrulli* survives on and in cucurbit seed from where it can spread to infect the cotyledons as they emerge from the seed coat. Seedlings germinating directly in the field or in glasshouses can be infested. Seedlings grown in warm and humid glasshouses are more susceptible to disease establishment and the rate of disease spread can be high. Infected seedlings may or may not show symptoms depending on the level of inoculum, virulence level of the strains, environmental conditions and host plant. Symptomless seedlings can be transplanted into the field. As plants grow in the field, the pathogen is disseminated by wind-driven rain, irrigation water, thundershowers or by mechanical means via farm workers and cultivation equipment. Cucurbit fruit can then become infected via natural openings or wounds, fruits may be infected internally through infection of female blossoms. Fruit can eventually rot in the field and their seeds, infested by contact with contaminated tissues, slip to the soil. The bacterium can overwinter in volunteer seedlings from seeds from infected fruits, infested rind or other infested crop residues as well as cucurbit weed hosts (especially wild citron, *Citrus medica*). Such material then acts as a local source of inoculum that may contribute to outbreaks in subsequent cucurbit crops. The disease cycle of bacterial fruit blotch of cucurbits, from the introduction of the pathogen with contaminated seed to the survival of the pathogen in the field, is illustrated in Latin & Hopkins (1995) and Walcott (2005). (Further details and references are provided in the datasheet).

**Figure ES1:** The disease cycle of bacterial fruit blotch of cucurbits. Steps 1 to 9 illustrate the cycle from the introduction of the pathogen with contaminated seed to the survival of the pathogen in the field.



Source: Figure reproduced from Latin & Hopkins (1995) with permission from R.X. Latin

- **Host range**

*Acidovorax citrulli* causes the disease bacterial fruit blotch on cucurbits, specifically on watermelon (*Citrullus lanatus*) and cantaloupe and honeydew melon (*Cucumis melo*) which are the most susceptible hosts. Other cultivated hosts include cucumber (*Cucumis sativus*), pumpkin (*Cucurbita moschata*) and squash (*Cucurbita pepo*). Citronmelon (*Citrullus lanatus* var. *citroides*) is a weed host. Recently betel vine (*Piper betle*: Piperaceae) was reported as a host in Taiwan.

Alternative hosts have been also suggested based on artificial inoculations: solanaceous plants/fruits and papaya fruits (Nascimento *et al.*, 2004).

- **Means of dispersal / spread**

Long distance dispersal of the pathogen is via contaminated seed or transplant seedlings moved in trade. Once *Acidovorax citrulli* has been introduced to cucurbit production local / natural spread occurs via wind-driven rain splash and mechanical means. Spread is most rapid on hot summer days with heavy afternoon thundershowers that includes blowing rains. In glasshouse systems overhead irrigation can spread the disease.

## Time period considered by this assessment

State the time horizon that you considered during this assessment. If climate change is taken into account, note the climate change scenarios considered.

Much new information has been published on this pathogen since 2009. More scientific papers are expected in the next few years. This assessment looks ahead up to five years and does not take climate change into account. It is suggested that the assessment be updated in the next few years to take into account any new information published in scientific literature.

## Geographic Distribution

North America: Present, USA in several, mostly southern, States, Canada

South America: Present, Brazil.

Europe: Outbreaks under eradication in Greece, Hungary and Italy.

Africa: Present, South Africa (unconfirmed). Sato (2009) lists bacterial fruit blotch in South Africa, no other details are provided. When contacted, Dr. Sato replied that the source of information is through personal communication with a colleague in S. Africa (Sato pers. comm., 2011) thus its presence in South Africa remains unconfirmed.

Asia: Present, China, Indonesia, India, Iran, Israel, Japan, South Korea, Taiwan, Thailand, Turkey.

Oceania: Present, Australia, Guam, Northern Mariana Is.

## Pathways

Three pathways are considered. The principle pathway which is most likely to facilitate spread is through use of contaminated cucurbit seed intended for planting; the other pathway likely to provide a route for entry into the EU is the use of contaminated seedlings, grown from contaminated seed lots, then transplanted and intended for cucurbit production; a third pathway considered, though much less likely to provide a route for pest establishment is through import of contaminated cucurbit fruit such as watermelons, other melons, pumpkins or cucumbers. However, the likelihood of bacteria transferring from contaminated fruit to growing plants is extremely small.

## Summary of risk elements

- Likelihood of entry and transfer (*summarise 2.08*)

*Acidovorax citrulli* is most likely to enter the EU via contaminated watermelon seed, or contaminated watermelon seedlings that have been grown from a batch of infected seed. Transfer and spread to other plants is almost certain via overhead irrigation, or wind-driven rain splash. Combining risk elements that contribute to the likelihood of pest entry and transfer using a BBN model (described elsewhere) suggests that there is a moderate (medium) to high likelihood that over the next five years the pathogen will enter and transfer to hosts in the EU. However, there is considerable uncertainty around this and there is some chance that the likelihood of entry and transfer could actually be lower or higher. Further study of how risk elements and pathways are combined in the BBN model is justified.

- Potential area occupied at time horizon (*summarise extent of spread, 3.02*)

In general, conditions must be “warm and wet” for *Acidovorax citrulli* to develop into bacterial fruit blotch of cucurbits. Conditions used to drive disease development in seed testing

suggests 55% relative humidity and temperatures between 24°C and 38°C are optimal although the specific range of temperature and humidity over which the disease can develop is unknown. Based on the locations of where the disease has been reported elsewhere, almost all of the watermelon growing area of the EU is likely to provide suitable conditions within which the disease could develop. Conditions in glasshouses that use overhead irrigation would also provide suitable conditions for the pathogen.

In a scenario with no specific phytosanitary measures being put in place, and with widespread trade of cucurbit propagation material within the EU, using seed or seedlings from Third Countries, it is possible that over the next five years the pathogen could be dispersed to many individual sites where it could establish. Such rapid dispersal is made possible via trade networks and “super connected” traders within a network. However, it is unknown how likely a super-connected member of a trade network would be to deal in infested seed or seedlings.

Reports from the USA indicate that disease incidence is localised and probably due to use of contaminated seed. Further spread from outbreak sites is generally localised and if good hygiene measures are used, the disease could be prevented from occurring the next year. Hence although without measures in the EU the pathogen could be spread to sites across the EU, the total area occupied is not likely to exceed more than 1/3 of the potential host area.

- Consequences (*summarise the combination of 3.03 and 3.04*)

Bacterial fruit blotch was first recognised as a significant pest in 1987 when the disease destroyed entire fields of watermelon crops on Guam and Tinian in the Mariana Islands. Since then, the disease has rapidly emerged as a serious pathogen of watermelon and melon crops in the USA and is a major threat to these industries around the world (Bahar & Burdman, 2010). In warm and wet conditions the consequences to the most susceptible crops such as watermelons and melons can be severe. For example, of 500ha affected in south western Indiana nearly 100 ha incurred losses approaching 90%. If similar losses occurred in the EU, the impact could be regarded as moderate to very high within the watermelon industry.

No serious environmental impacts are expected to be caused by the pathogen.


- Pest risk (*summarise 3.06, the combination of likelihood of entry & transfer 2.0, with potential impact, 3.05*).

*Acidovorax citrulli* is a pathogen of cucurbits, especially damaging to watermelons, which has spread its geographic distribution in recent years and when environmental conditions are suitable, major crop loss can result.

An experimental risk assessment method and BBN model (described elsewhere) was used to systematically combine elements contributing to risk. Model outputs are presented in a series of bar charts. Figure ES1 shows the final model output which combines scores for likelihood of entry with potential impact to give an overall pest risk profile. The profile shows much uncertainty with some likelihood of risk being minimal (i.e. very low) up to major (very high). Most likely the pest risk is actually medium to high. Taking into account the major impacts reported where outbreaks have occurred, but which are normally only reported in a

small area, the overall rating of medium pest risk presented to the EU by *Acidovorax citrulli* seems a reasonable output for the model and is in line with assessors opinion.

**Figure ES 1:** Graphical representation of Pest Risk (3.06), combining overall potential for pest entry and transfer (2.08) with potential impact (3.05).



### Uncertainties

Summarise major uncertainties and report on what happens if those risk elements that are most uncertain are changed (e.g. become less uncertain) – how does the overall risk profile of expected risk change? Describe changes in the bar chart showing expected risk if uncertainty is changed.

Three pathways were considered (cucurbit seed for planting, cucurbit seedlings for planting and cucurbit fruit for consumption). The assessment focussed on watermelon seed, seedlings and fruit from countries where the disease has been reported. A critical uncertainty in all of the pathways was estimating the likelihood of the pathogen being associated with the commodity at the start of the pathway. The pathway of seedlings had uncertainty around how likely infected seedlings would be detected and removed from the pathway during routine quality checks. Further research on the detailed environmental conditions required by the pathogen to develop symptoms would better inform the likelihood of establishment in the risk assessment area and also reduce the uncertainty around identifying the endangered area.

### Conclusion

The bacterial plant pathogen *Acidovorax citrulli* has the characteristics of a quarantine pest for the EU.

**Keywords:** bacterial fruit blotch of cucurbits, *Acidovorax citrulli*, pest risk assessment

## Stage 1 – Initiation

### 1.1 Background and Initiation

Provide the background and terms of references as provided by the originator of the risk assessment request (European Commission, European Parliament, Member States, or EFSA)

The purpose of this assessment is to evaluate the plant health risk of *Acidovorax citrulli* within the framework of EFSA project CFP/EFSA/PLH/2009/01 (Prima phacie).

The terms of reference are described in EFSA call CFP/EFSA/PLH/2009/01, Pest risk assessment for the European Community plant health: A comparative approach with case studies (EFSA, 2009). The text in Section 1.4 of the call, “Structure and essential requirements of the proposal”, pages 7-9, provide the terms of reference e.g. that a systematic review of risk assessment methodologies, with emphasis on quantitative and semi-quantitative approaches, used in pest risk assessment to analyse and predict the likelihood of entry, establishment and spread, the potential negative consequences, the overall risk characterisation and the associated level of uncertainties be assessed, together with a systematic review of the methods used to assess the effectiveness of management options in reducing the risk of introduction and/or spread. The quantification of economic losses in monetary values and the assessment of potential effects on export markets, employment and tourism were not to be included.

#### Initiation Point

This assessment was initiated as a case study pest to be examined within EFSA project CFP/EFSA/PLH/2009/01 (Prima phacie). *Acidovorax citrulli* had been selected as a case study pest because it satisfied a number of criteria needed to provide a range of contrasting pest examples for consideration in the project.

### 1.2 Identification of the risk assessment area

The risk assessment area is the 27 Member States of the EU with the focus on the continental European area, specifically excluding the ultra-peripheral regions, i.e. the French overseas departments, Spanish Canary Isles and Portuguese Azores and Madeira.

### 1.3 Available pertinent regulatory information

#### (i) Previous risk assessment or pest risk analysis?

A pest risk analysis was performed by Üstün Nursen from the Plant Protection Research Institute in Turkey (Nursen, 2008) following the finding of an emerging disease referred to as “watermelon fruit blotch” detected in Turkey in 1995. The analysis considered Turkey as the PRA area.

#### (ii) Available Pest Fact Sheets/ Pest Alerts etc.

A pest datasheet for *Acidovorax citrulli* was composed and submitted as Annex 1d to the first Prima phacie interim report (July, 2010). The datasheet has subsequently been updated with additional information, for example regarding its name (*A. citrulli* rather than *A. avenae* ssp. *citrulli*), and its geographic distribution.

The CABI Crop Protection Compendium also has a datasheet for this organism, using the name *Acidovorax avenae* ssp. *citrulli* (last updated by CABI 08 April 2011 (CABI, 2011)). There is a short datasheet by EPPO on the pest (EPPO, 2009a).

### (iii) Current regulatory status

What is the pest's status in the Plant Health Directive (Council Directive 2000/29/EC<sup>4</sup>) ?

*Acidovorax citrulli* (or its synonyms) is not specifically mentioned in the EC Plant Health Directive.

*Acidovorax citrulli* is not specifically mentioned in the EC Marketing Directive 92/33/EEC of 28<sup>th</sup> April 1992, although there is a requirement that vegetable propagating and planting material (other than seed) of *Cucumis melo* (melon), *Cucumis sativus* (gerkin), *Cucurbita pepo* (courgette), *Cucurbita maxima* (gourd) be substantially pest free.

### (iv) What is the pest's status in the European and Mediterranean Plant Protection Organisation (EPPO)? (put tick (✓) in box if relevant) ([www.eppo.org](http://www.eppo.org))

EPPO List: A1 regulated pest list  A2 regulated pest list  Action list  Alert list

*Acidovorax citrulli* was added to the EPPO Alert List in July 2009 after the EPPO Panel on bacterial diseases considered that it was a possible seed-transmitted threat to cucurbit crops in particular melon and watermelon, following reports of its spread in other parts of the world (EPPO, 2009b).

### 1.4 Strategy of data searching (identity of databases, data banks and information systems, key search terms and strategies applied, and the time period covered should be provided)

Information searches were performed consulting several sources such as:

- Abstracting databases: e.g. AGRICOLA, CAB Abstracts, ISI Web of Knowledge
- Internet search machines: Google Scholar
- EPPO information systems: e.g. EPPO reporting service, EPPO PQR v 5.0.1500
- Information from Member States on issues related to host distribution at a national level was acquired via a questionnaire prepared in the framework of the Prima Phacie project and distributed by EFSA to all NPPOs.

---


<sup>4</sup> <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:32000L0029:EN:NOT>

### 1.5 Time period considered in this assessment

(Likelihoods of future events are only meaningful when a time period is specified. Choices that might be considered include: 3 months (short term, e.g. following discovery of an outbreak or other new potential threat), 5 years (medium term), or 30 years (long term). Choose a time scale and briefly explain why such a horizon was selected. – note whether climate change was taken into account. If climate change is taken into account, note the climate change scenarios considered.

This assessment takes a medium term view, looking up to five years into the future. The chart below draws on the list of references cited in the Prima phacie datasheet for *Acidovorax citrulli*. The chart shows the number of publications that are cited in the datasheet against the year the citation was published, each year since 1965. Approximately one third of all references cited have been published in the last 4 years (i.e. since 2008). The trend shows that the rate of publications relating to this pest has increased in recent years. Anticipating that such a trend will continue for the next few years at least, it can be expected that new information will be published to further inform any future risk assessment. In anticipation of further information to emerge in the next few years, it seems unwise to speculate very much further than a few years into the future and taking a much longer term view is not appropriate. If the number of publications were declining or stabilising, it could suggest that sufficient information about the biology of the pest was available to permit a longer term time horizon to be considered perhaps.

**Figure 1.5:** Number of references cited in the datasheet for *Acidovorax citrulli* by year of publication



## 1.6 Introductions or interceptions (reported from EU or elsewhere)

Provide information on interceptions or reports of pest spread and note whether the pest has a history of increasing its area of distribution.

The disease, bacterial fruit blotch of cucurbits, was first reported in the USA in areas of commercial watermelon production in 1989 (Hopkins, 1989; Latin & Rane, 1990). However, reports of symptoms and damage to watermelons probably caused by what is now called *A. citrulli* date back to Georgia and Florida in the 1960s (Latin & Hopkins, 1995) although the identity of the causal agent could not be ascertained at that time. Only in 1990 was it concluded that the disease was seed borne (Wall *et al.*, 1990) by which time the disease was being reported in more US states and from other countries around the world. Table 1.6 shows a time line of published records reporting the first occurrence of bacterial fruit blotch of cucurbits or *A. citrulli* in US States and countries since 1965.

Table 1.6: Time line of reporting new and first occurrence of bacterial fruit blotch of cucurbits.		
Year	Place of occurrence	Reference
1965	Georgia (USA)	Webb & Goth (1965)
1967	Florida (USA)	Crall & Schenck (1969)
1978	Queensland (Australia)	Queensland DPI (1978)
1986	Australia	Somodi <i>et al.</i> (1991)
1987	Mariana Islands (Guam, Tinian)	Wall & Santos (1988)
1989	USA (Iowa, Delaware, Maryland)	Latin & Hopkins (1995)
1989	USA (Indiana, Alabama)	Latin & Rane (1990); Sikora (2004)
1989	USA (Delaware, Florida)	Evans & Mulrooney (1991); Somodi <i>et al.</i> (1991)
1989	USA (North Carolina)	Bailey 1996
1991	USA (Oklahoma)	Jacobs <i>et al.</i> (1992)
1992	USA (Georgia)	Hopkins <i>et al.</i> (2000)
1993		
1993	USA (Texas)	Black <i>et al.</i> (1994)
1994	USA (Missouri)	Jett <i>et al.</i> (2002)
1995		
1995	Turkey	Demir (1996)
1996	USA (Oregon)	Hamm <i>et al.</i> (1997)
1997	Nicaragua*	Munoz & Monterroso (2002);
1998	China (Hainan)	Feng <i>et al.</i> (2009)
1991	Brazil	Assis <i>et al.</i> (1999); Oliveira <i>et al.</i> (2003) Robbs <i>et al.</i> Fitopatol. Bras. 16:48, 1991)
1998	Japan	Shirakawa <i>et al.</i> (2000)
2001	China (Xinjiang, Neimenggu)	Zhao <i>et al.</i> (2001)
2001	USA (Illinois);	Babadoost & Pataky (2002);
2002	Costa Rica	Mora-Umana & Araya (2002)
2003		
2004		
2005	China (Fujian)	Cai <i>et al.</i> (2005)
2006		
2004	Iran	Harighi (2007)
2007	Hungary	Palkovics <i>et al.</i> (2008)
2005	Greece;	Holeva <i>et al.</i> (2009);
2009	Italy *	EPPO (2010)
2008	Taiwan	Deng <i>et al.</i> (2010)
2011	USA (Virginia, Delmarva Peninsula, Ohio, Kentucky)	Langston (2011) (radio interview)

\* under eradication or subsequently declared eradicated

Note that the disease has not necessarily persisted in all regions where it has been reported. For example the disease occurred in Iowa in 1989 and up to 1995 had not been seen again; similarly outbreaks in Delaware and Maryland in 1989 have not recurred, supporting the suggestion that in northern US States of America the disease relies on the pathogen being carried and introduced on infected planting material (Latin & Hopkins, 1995) suggesting that the pathogen may be unable to persist outdoors in northern US climates. However, in southeastern US perennation of the pathogen may be possible as cucurbit weeds that survive in ditch banks and wooded areas may harbour the bacteria (Latin & Hopkins, 1995)

The finding of bacterial fruit blotch in Nicaragua followed the import of watermelon seed from Costa Rica in 1997 (Muñoz and Monterroso, 2002). In Turkey, the disease was first found in 1995 in the Edirne province on watermelon crops (Demir, 1996) and in 2005 it was in Adana Province (Mirik *et al.*, 2006). In Hungary, the disease was observed in watermelons in July 2007 during a hot summer period. Although the source of infection could not be determined, watermelon transplants had been imported from Turkey (Palkovics, 2008).

In Greece, the disease was found in July 2005 and September 2006, on mature watermelon fruits originating from the areas of Chryssoupoli (Macedonia, northern Greece) and Vagia (central Greece), respectively. Subsequently, the disease was found in May 2008, on young grafted watermelon plants from the area of Varda (Peloponnese, southern Greece) (Holeva *et al.*, 2010). Since then, there has been no record of any other incidence of the disease.

US watermelon seeds contaminated with *A. citrulli* were intercepted on a number of occasions by Israel between 1992 and 1994 (Assouline *et al.*, 1996). The only report of diseased watermelon fruit being intercepted is from the Mariana Islands when diseased fruit moving from Tinian to Guam was intercepted by phytosanitary inspectors (Wall *et al.*, 1990). Transfer from infected fruit to a growing crop seems unlikely unless the fruit is disposed of carelessly, very close to a site of production.

From the summary above, it is clear that there is recent history of reporting new occurrences of *A. citrulli* and bacterial fruit blotch. Most frequently it is suggested that the pathogen is spread via contaminated seed (e.g. interceptions by Israel and its entry into Nicaragua from Costa Rica via seed) or seedlings (e.g. occurrence in Hungary after importing seedlings from Turkey).

## Stage 2 - Pest Risk Assessment

(Outline approach)

This method for pest risk assessment involves first evaluating the likelihood of pest entry and transfer to a host within the risk assessment area. Likelihood of entry is assessed by considering five factors:

- (i) likelihood of association with commodity on the pathway at origin,
- (ii) pest survival during post harvest treatment,
- (iii) pest survival during storage and transport,
- (iv) pest survival during current phytosanitary procedures, and
- (v) the quantity of commodity imported.

The likelihood that sufficient numbers of pest will transfer from a pathway to a suitable host in order to initiate a new population is then considered. The combined likelihoods of entry and transfer via each pathway are then combined before likelihood of establishment is taken into account. Assessors then move onto assess consequences of establishment.

Each risk element or sub-element is divided into five categories. Assessors review data / evidence and allocate % likelihood to appropriate categories, either selecting a single category or spreading their judgment between categories. Guidance is provided to interpret the categories in order to provide some consistency.

Overall potential impact is determined via use of BBN software based on matrices that combine consequences of establishment with establishment potential **given entry and transfer**. Likelihood of entry and transfer is then combined with potential impact using the BBN software to estimate pest risk.

### Acknowledgement

Method 4b has largely adopted questions from the USDA pathway initiated pest risk assessment method (USDA, 2000). However, the arrangement and structure of questions has been revised by Prima phacie so that the method is more aligned with EFSA needs.

### Reference

USDA Guidelines for pathway initiated pest risk assessments v 5.02 (2000).

## Likelihood of pest entry and transfer to a host

### 2.0 List and describe the pathways for pest entry into the risk assessment area

A pathway is “any means that allows the entry or spread of a pest” (ISPM No. 5, IPPC, 2007). Remember to consider potential pathways that are closed due to existing phytosanitary measures but which could be opened if the phytosanitary measures were changed.

Entry is “Movement of a pest into an area where it is not yet present, or present but not widely distributed and being officially controlled” (ISPM No. 5, IPPC, 2007).

Copy 2.1a to 2.7a for each pathway, and give responses for each individual pathway. Label questions 2.1b for the second pathway assessed, 2.1c for the third pathway etc.

Pathway	Pathway name	Summary description of pathway
a.	<b>Seeds of cucurbits.</b>	Import of seeds of cucurbits, especially watermelon, from countries where <i>Acidovorax citrulli</i> occurs.
b.	<b>Seedlings (transplants) of cucurbits</b>	Import of seedlings of cucurbits for fruit production from countries where <i>Acidovorax citrulli</i> occurs.
c.	<b>Fruits of cucurbits</b>	Import of cucurbit fruit for consumer consumption from countries where <i>Acidovorax citrulli</i> occurs.

Note that on occasions risk assessments are initiated by a review of phytosanitary policy, e.g. for pests already present and perhaps widespread in an area and which therefore may no longer be suited to being dealt with using official phytosanitary measures. In such a situation, the likelihood of pest introduction is relevant only for limited parts of the area where the pest does not occur, or where it is not widely distributed and remains under official control. Assessment of pest entry could then be restricted to such areas.

### Uncertainties (regarding pathways)

Following EFSA Guidance (EFSA, 2010), to ensure transparency in risk assessment, uncertainties should be identified, characterized and documented within all risk assessments. Identifying uncertainties can show not only which aspects of an assessment are uncertain but the degree of uncertainty and can help identify where further work could usefully reduce uncertainty.

Describe the uncertainties regarding the identification of pathways.

There is a lack of detail regarding precise sources of material on each pathway, the time of year material is moved, volumes, routes and modes of transport used. Assumptions had to be made about the environmental conditions in which commodities are transported.

In terms of additional pathways, there is one report from Israel suggesting that *A. citrulli* may be transmitted by solanaceous seeds – the pathogen has been isolated from tomato and aubergine seeds imported to Israel (Assouline *et al.*, 1997). However, this pathway was not examined due to low likelihood of transfer to a host. Nevertheless with more time such a potential pathway would be worth assessing, particularly if *A. citrulli* expands the range of hosts on which disease is expressed.

**Pathway 1: Seeds of cucurbits** (Seeds of cucurbits, especially watermelon, from countries where *Acidovorax citrulli* occurs)

**2.01a Likelihood of the pest being associated, spatially and temporally, with the pathway at origin**

(There must be some likelihood of association otherwise there is no pathway).

Take into account pre-harvest cultivation and husbandry practices such as existing pest management measures, choice of cultivar, and applications of plant protection products. If phytosanitary measures (i.e. statutory risk reduction measures) are already in place against this or other pests at origin, specify whether these are being taken into account or not.

**Information / evidence:** *Provide reasoning then give judgment*

The EU does not provide data on exports and imports of specific vegetable seed hence there is very little data available about the EU import and trade of cucurbit seeds. However, anticipating the lack of statistics, the Prima phacie case study consortium developed a questionnaire for EU MS to address some anticipated information gaps (Annex 1). Part of the questionnaire was designed to determine which EU MS import cucurbit seed and from where such seed is obtained. A summary of responses is shown in Table 2.01a.

<b>Table 2.01a: Third country sources of cucurbit seed imported by EU MS</b>						
TC seed source	NL	DE	IT	PT	FI	CZ
USA *	x	x	x	x	x	
Israel *	x	x	x	x		
Thailand *	x	x				x
China *	x		x			
South Korea *				x		
India *	x					
Japan *	x					
Taiwan *	x					
Chile	x					
Indonesia	x					
Peru	x					
Morocco	x					
Tanzania	x					
New Zealand		x				

Source: Prima phacie questionnaire to EFSA PLH Network

Key: x = cucurbit seed has been imported from this Third Country

\* = *Acidovorax citrulli* present in this county

Table 2.01a shows that at least six EU MS have sourced cucurbit seed from countries where *A. citrulli* occurs, with USA and Israel being the most common sources. Other EU MS may also use seed from these and other Third Countries but seed can be imported and repackaged within the EU and the origin of seed can then be lost. The same is true for seed imported from a Third Country via another Third Country, e.g. import of Indian seed to Israel via California (USA) (Assouline *et al.*, 1997).

It can be concluded that the pathway of seeds of cucurbits imported to the EU from countries where *Acidovorax citrulli* occurs is a real pathway.

Regarding likelihood of association at origin, recent studies have shown that infected fruit may contain *A. citrulli* infected seeds regardless of whether the fruit shows symptoms or not (Walcott et al., 2003; Lessl et al., 2007; Bahar et al., 2009). This causes problems for seed companies aiming to produce *A. citrulli* free seeds and means that it is very difficult to certify that seed may be pest free, even if good field sanitation practices, such as removal of fruit debris, volunteers and wild hosts, have been in place at the place of production. No resistant cultivars have been identified, though some tolerance has been found in watermelons and melons (Hopkins & Thompson, 2002; Bahar *et al.*, 2009). There are no figures available for percentage infection amongst the seed being taken for growing stock, but the fact that money and effort is being put into post harvest treatments of seed suggests that sourcing seed from countries where *A. citrulli* occurs does present a significant likelihood of association.

On the other hand, there have been only a few incidents of bacterial fruit blotch outbreaks on cucurbits in the EU despite a history of importing cucurbit seed from countries where the disease occurs. Nevertheless, an outbreak of bacterial fruit blotch in Nicaragua was linked to imported seed (EPPO, 2009a) as was an outbreak in Turkey (Demir, 1986; Nursen, 2008) and in Indiana in USA (Latin & Hopkins, 1995) so it is possible that there will be some association at origin.

The interception of infested seeds, e.g. i) watermelon seeds originating in USA have been intercepted several times in Israel between 1992 and 1994; ii) tomato seed from India intercepted in Israel, indicates that association of the pathogen with seed is possible (Assouline *et al.*, 1997).

In addition, Zhao *et al.* (2009) reported that preliminary seed health testing in China revealed that up to 70% of the Hami melon seed lots in some areas was contaminated with the pathogen.

#### **Uncertainties regarding likelihood of the pest being associated with the pathway at origin**

Data on the specific area of origin of cucurbit seed within source countries is unavailable. It is unknown if any seed is likely to be sourced from areas where outbreaks have occurred.

## Conclusions

<b>2.01a: Likelihood of association with the pathway at origin</b>			
<b>Rating</b>	<b>Description</b> (likelihood of association is ....)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the commodity are likely to be contaminated / infested)	Expect higher association given that outbreaks have been found associated with seed	-
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> are likely to be contaminated / infested)	Expect higher association given that outbreaks have been found associated with seed	-
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the commodity are likely to be contaminated / infested)	Very difficult to tell if seed is pest free – fruit may show no symptoms.	80%
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the commodity are likely to be contaminated / infested)	Although few consignments are imported (see 2.05a) a few outbreaks have occurred so expect some association	20%
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the commodity are likely to be contaminated / infested)	If association was at this level or higher more interceptions of outbreaks would be expected	-
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence.

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

*The assessment of pest entry and transfer (2.01 -2.06) is based on considering “lots”.  
However, if an alternative unit would be more appropriate then describe the chosen unit and ordinal scale using five categories.*

## **2.02a Likelihood of surviving postharvest treatments / measures (before pest entry into risk assessment area)**

*Given that a proportion of lots/ consignments may be infested / contaminated, consider the proportion of infested/contaminated lots that are likely to remain infested/contaminated after any manipulation, handling or specific phytosanitary treatment to which the commodity is subjected. Examples of postharvest treatments include culling, washing, chemical treatment and cold storage. If there are no post harvest treatments the likelihood of survival should probably be considered “very high”.*

*If post-harvest phytosanitary measures (i.e. statutory risk reduction measures) are already in place, specify whether these are being taken into account or not.*

### **Information / evidence:** *Provide reasoning then give judgment*

A range of seed treatments (e.g. heat, NaOCl, HCl, Peroxyacetic acid and fermentation with CaOCl<sub>2</sub>) have been investigated to try and eliminate *Acidovorax citrulli* but, although there has been success in reducing incidence in infected seeds, none have been found to completely eradicate the pathogen, without reducing seed quality (Rane & Latin, 1992; Hopkins *et al.*, 2003). Feng *et al.* (2007), Hopkins *et al.* (1996), Sowell & Schaad (1979), Wall (1989) and Assouline *et al.* (1997) reported detection of the pathogen from tomato seeds having a certificate stating to have been treated by hot water at 50°C for 60 min. Rane & Latin (1992) reported the recovery of the pathogen from both the seed coats and the embryos of seeds from contaminated fruits, which indicates that seeds are both internally and externally contaminated. Similar observations were made by Neto *et al.* (2006). Thus, the inability of the various treatments to eradicate the pathogen from the seed could be explained.

It has also been observed that there are openings in the watermelon seed coat at the hilum region (visible with scanning electron microscopy) through which microorganisms could enter during the seed extraction process. This mode of seed transmission has been found in other bacterial pathogens e.g. *Xanthomonas axonopodis* pv. *phaseoli* and *Pseudomonas syringae* pv. *phaseolicola* (Rane & Latin, 1992).

### **Uncertainties regarding likelihood of the pest surviving postharvest treatments/ measures (before pest entry into the risk assessment area)**

No seed treatment measures are entirely effective and the proportion of seeds that remain infested post treatment is unknown.

### Conclusions

Assuming no specific measures are in place against *A. citrulli* the likelihood that infested seed remains infested is very high.

<b>2.02a: Likelihood that an infested/contaminated commodity remains infested/contaminated after existing postharvest treatments/measures</b>			
<b>Rating</b>	<b>Description</b> (likelihood of remaining infested/contaminated / pest survival is ..)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Without specific measures applied there is no reason to expect contaminated seed to become pest free	100%
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

### **2.03a Likelihood of surviving storage and transport**

Given that a proportion of lots/ consignments may still be infested / contaminated, estimate the proportion of lots that are likely to remain infested/contaminated because the pest can survive storage and transport; consider speed and conditions of transport and duration of the life cycle of the pest in relation to time in storage and transport, commercial procedures (e.g. refrigeration) applied to consignments in the country of origin, during shipping, and in the country of destination, that could affect the likelihood of pest survival. Take into account previous live interceptions on this or similar pathways (see 1.6).

If phytosanitary measures (i.e. statutory risk reduction measures) are already in place which act on the likelihood of pest survival during storage and transport, specify whether these are being taken into account or not.

#### **Information / evidence:** *Provide reasoning then give judgment*

Contaminated seed is very likely to remain contaminated during transport. Survival in storage of up to at least 26 months is also very likely.

Hopkins *et al.* (1996) showed that the level of disease transmission was not reduced after contaminated seed had been stored at 12°C for 12 months. Shirakawa *et al.* (2003) conducted trials on the survival of *A. citrulli* on artificially contaminated seeds at fixed temperatures between 4 and 30°C over 26 months. The bacterium was able to survive at all temperatures. Long term storage of seed, if stored in conditions suitable for seed survival, i.e. in a cool and dry environment, is not likely to eliminate any contaminated seed (Hopkins *et al.*, 1996). Furthermore, reports that the bacterium can be transmitted to seedlings from infested seeds stored for more than 38 years prompted Dutta & Walcott (2010) to compare the ability of *A. citrulli* to survive on seeds with that of *Xanthomonas campestris* pv. *campestris*, *Ralstonia solanacearum* and *Pantoea stewartii* subsp. *stewartii*. They found that the first three bacterial species are more tolerant to desiccation than *Pantoea stewartii* subsp. *stewartii*. They suggested that the ability of *A. citrulli* to survive for 38 years in stored seed is due to the location of the bacterium in the seed rather than some unique characteristic of the bacterium.

#### **Uncertainties regarding likelihood of the pest surviving storage and transport**

It is unknown how long cucurbit seed could be kept in storage. Storage of over 26 months seems unlikely on a commercial scale. However, survival has been recorded for at least 26 months.

### Conclusions

Assuming no specific measures are in place against *A. citrulli* the likelihood that infested seed remains infested is very high.

<b>2.03a: Likelihood of surviving storage and transport</b>			
<b>Rating</b>	<b>Description</b> (likelihood of remaining contaminated / pest survival is ...)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Without specific measures applied there is no reason to expect contaminated seed to become pest free	100%
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

## 2.04a Likelihood of pest surviving current phytosanitary procedures at the point of entry or elsewhere in the risk assessment area

Given that a proportion of lots may still be infested / contaminated, estimate the proportion of lots that are likely to remain infested/contaminated because the pest survives existing phytosanitary procedures e.g. it is not detected at entry and/ or it can survive any existing phytosanitary procedures within the pest risk assessment area. Take into account the intensity of sampling and inspection and ease of detecting and distinguishing the pest from other organisms.

### Information / evidence: *Provide reasoning then give judgment*

Pathogen free seeds or transplants can be achieved through a thorough inspection of the plant material before introduction to the growing area (glasshouse or field). However, this is not a regulated pest and the Plant Health Directive (2000/29 EC) makes no requirement to conduct phytosanitary checks on cucurbit seeds imported to the EU. Tests that could be used are very expensive in relation to the cost of seed. In the USA seed suppliers do conduct tests but do not give guarantees that marketed seed is entirely free from *A. citrulli* (Latin & Hopkins, 1995). In Israel, where the pathogen is regulated, there have been reports of interceptions of infested seeds (Assouline *et al.*, 1997).

### Uncertainties regarding likelihood of the pest surviving current phytosanitary procedures

It is unknown what proportion of imported seed is checked and tested voluntarily.

### Conclusions

No phytosanitary procedures are currently imposed, therefore highly likely to survive.

2.04a: Likelihood of pest surviving current phytosanitary procedures			
Rating	Description (likelihood of remaining contaminated/ pest survival is ...)	Justification summary	Probability Assignment <sup>1</sup>
Very Low	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
Low	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
Medium	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
High	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
Very High	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Checks are possible, but not required	100%
	Check sum =		100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup> **Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

## 2.05a Quantity of commodity imported annually

*Quantity of commodity imported annually: The likelihood that a pest will enter depends on the amount of the potentially-infested commodity that is imported. For qualitative pest risk assessments, the amount of commodity imported is estimated in units of tonnes, or other metric such as standard 40 foot long shipping containers.*

*If the quantity of commodity imported is better described using alternative units, such as the number of plants for planting, assessors should devise a 5 level scale and provide some reasoning to support use of the scale.*

### **Information / evidence:** *Provide reasoning then give judgment*

The EU does not provide data on exports and imports of specific vegetable seed hence there is very little data available about the EU import and trade of cucurbit seeds. However, some countries such as the US, India and Chile break their vegetable seed exports down by crop and country.

For tariff and tax reasons customs authorities record the quantity of many commodities and goods that are traded internationally. Such commodities can be described using a six digit number within a system called the Harmonised System (HS). The HS is a high-level international nomenclature owned and developed by the World Customs Organisation. For example, within the HS, 060240 refers to “rose plants, whether grafted or not”. However, within the EU a more detailed code system of 8 or 10 digits is used. This divides the HS codes into ever more detailed descriptions of goods. Thus whilst 06024 refers to rose plants, whether grafted or not, 06020411 refers to roses with a stock diameter of up to 10mm, whilst 06024019 refers to roses with a stock diameter of more than 10mm. The EU system is called the Combined Nomenclature (CN).

Many fruit and vegetable commodities are described within the HS and CN systems. Unfortunately the trade in seeds of plants are not well described within the HS or CN systems. Nevertheless, the International Seed Federation (ISF)<sup>5</sup> does provide basic statistics about fruit, flower and field crop seed exports and imports. It is not known whether seeds of watermelons, melons, cucumbers, pumpkins and other cucurbits are regarded as vegetable seeds or field crops. It is assumed that field crops are mainly arable crops, e.g. cereals, and that cucurbits are regarded as vegetables. Thus Table 2.05a 1 show EU imports of vegetable seeds. In 2010 almost 100,000 tonnes of vegetable seeds were imported by EU countries. However, only a small proportion of the seeds are likely to be cucurbit seeds.

EU MS do import cucurbit seed from a number of countries where *A. citrulli* occurs (see Table 2.01a previous). Table 2.05a 2 shows the quantity of all vegetable seed exported by countries where *A. citrulli* occurs, or where the status of the pathogen is unconfirmed (South Africa) or where outbreaks of the disease have been reported. As with EU imports, only a small proportion of the export seeds are assumed to be cucurbit seeds. A relationship between EU imports and countries exports cannot be made using this data. However, there is undoubtedly an international trade in cucurbit seeds as evidenced by responses to the Prima phacie questionnaire (Table 2.01a).

---

<sup>5</sup> <http://www.worldseed.org/isf/home.html>

Table 2.05a 1: EU imports of vegetable seeds for planting, 2010 (Tonnes)			
EU MS	Tonnes	EU MS	Tonnes
Belgium	21,434	Ireland	764
Netherlands	18,672	Sweden	700
Austria	16,587	Lithuania	577
Italy	9,976	Slovakia	400
Germany	5,766	Czech Republic	379
UK	5,654	Denmark	343
France	4,450	Cyprus	275
Spain	2,790	Slovenia	256
Hungary	2,541	Bulgaria	215
Greece	2,342	Malta	200
Poland	2,220	Finland	180
Romania	1,504	Latvia	100
Portugal	770	Estonia	50
Sum			99,145

Source: [http://www.worldseed.org/isf/seed\\_statistics.html](http://www.worldseed.org/isf/seed_statistics.html)

Note that only a fraction of the quantity of seed imported into the EU is likely to be cucurbit seed.

Table 2.05a 2: Exports of vegetable seeds for planting from countries where <i>A. citrulli</i> occurs or where its status is unconfirmed, 2010.		
Country	Quantity (tonnes)	Value (US \$'000,000)
USA *	21,603	485
Italy (*)	10,453	106
Israel *	unknown	106
China *	5,742	88
Hungary *	2,039	17
Japan *	1,406	98
South Africa (**?)	1,253	17
Australia *	1,036	22
Turkey *	644	13
Korea, Republic of *	397	20
China, Taiwan *	unknown	11
Costa Rica *	0	10
Greece (*)	312	1
	44,885 + unknowns	994

Source: [http://www.worldseed.org/isf/seed\\_statistics.html](http://www.worldseed.org/isf/seed_statistics.html)

(\*) = outbreaks    \* = present    (\*\*?) = unconfirmed

Note that only a fraction of the quantity of seed exported by the countries in Table 2.05a 2 is likely to be cucurbit seed. Note also that the table shows total exports, not all of which would be destined for the EU.

Seed import statistics available for the USA show that over the past 20 years, the majority of imported watermelon seeds for planting in the US have come from Egypt or Israel. US data since 2005 is shown in Table 2.05b 3, below.

	2005	2006	2007	2008	2009	2010	6 year mean	% of 6 year mean
Egypt	137.5	78.0	43.3	61.7	74.6	62.0	76.2	29.6
Israel *	183.4	114.5	32.3	71.6	7.9	5.3	69.2	26.9
China *	18.3	23.3	19.6	31.0	40.9	16.5	24.9	9.7
Chile	26.0	37.0	28.0	10.8	10.2	27.8	23.3	9.1
Taiwan *	8.2	14.0	6.6	16.1	24.3	9.3	13.1	5.1
Peru	7.8	7.0	6.9	6.3	6.1	11.1	7.5	2.9
Mexico	16.5	3.6	1.3	1.8	5.6	4.0	5.5	2.1
Thailand	3.6	3.7	4.0	7.9	3.3	3.5	4.3	1.7
Japan *	0.1	0.0	0.5	0.8	1.0	1.0	0.5	0.2
All others	41.2	55.0	28.9	9.2	29.0	32.2	32.6	12.7
Sum	401.4	281.2	142.3	208.1	173.9	140.5	257.1	100.0

Source: Extracted from Table 19, Watermelon seed US Imports by country 1980 – 2010. Available at USDA Economics, Statistics and Market Information System <http://usda.mannlib.cornell.edu/MannUsda/viewDocumentInfo.do?documentID=1399>

Since 2005, on average, the USA has imported just over 100 tonnes of watermelon seed from countries where *A. citrulli* occurs (Israel, China, Taiwan, Japan). Although no data could be found regarding the quantity of cucurbit seed imported by all EU MS, the Dutch Plant Protection service was able to provide some information (Table 2.05a 4) which shows that over 85% of cucurbit seed imported by the Netherlands was sourced from countries where *A. citrulli* is present.

Year	2008	2009	2010	3 year mean	Mean annual %
China *	16,209	11,065	21,608	16,294	84.4
Thailand *	209	939		383	2.0
USA *	51	75	122	83	0.4
Israel *	1	192	35	76	0.4
Japan *	100			33	0.2
Taiwan *	37			12	0.1
<i>Sub total</i>	<i>16,607</i>	<i>12,271</i>	<i>21,765</i>	<i>16,881</i>	<i>87.5</i>
Chile	640	2,294	2,106	1,680	8.7
Peru	264	1,135	485	628	3.3
Morocco	155	186	16	119	0.6
Tanzania	5	18		8	0
Indonesia	1			0	0
<i>Sub total</i>	<i>1,065</i>	<i>3,633</i>	<i>2,607</i>	<i>2,435</i>	<i>12.6</i>
sum	17,672	15,904	24,372	19,316	100

Source: NL Plant Protection Service, pers. comm.. \* = *Acidovorax citrulli* present

Table 2.05a 4 shows that between 2008 and 2010 the Netherlands imported between 11.3 tonnes (2009) and 21.8 tonnes (2010) of cucurbit seed from countries where *A. citrulli* occurs.

The Royal Botanic Gardens at Kew has compiled a database of seed biological trait data from published and unpublished sources (<http://data.kew.org/sid/>). The primary purpose is as an internal source of a variety of seed biological information; for use in large scale analysis and decision support for seed conservation operations. The database provides the 1,000 seed weight for many plant species and varieties, including watermelons, melons and cucumbers (Table 2.05a 5). The 1,000 seed weight for watermelons, melons and cucumbers was used to estimate the number of cucurbit plants that could potentially develop from 10 tonnes of imported seed, or 25 tonnes of imported seed. Such quantities are within the order of trade occurring in recent years between NL and Third Countries.

<b>Table 2.05a 5:</b> Estimating the number of cucurbit seeds imported by NL based on 1,000 seed weight				
Crop plant	1000 seed weight (g)	Approximate number of seeds in 1 kg	Approximate number of seeds (millions)	
			in 10 tonnes	25 tonnes
watermelon	85.0	12,000	120	300
cucumber	16.6	60,000	600	1,500
melon	7.2	140,000	1,400	3,500

In recent years, in the order of 10 to 25 tonnes of cucurbit seed have been imported by the Netherlands from countries where *A. citrulli* occurs (Table 2.05a 4). This equates to approximately between 120 million seeds of watermelons (if all cucurbit seed imports were watermelon) to 3,500 million seeds of melon (if all cucurbit seed imports were other melons). Of all cucurbit seed imported, only a proportion of it would have been watermelon seed, the suspected primary source for introduction of *A. citrulli*. Assuming that across the whole of the EU between 2 and 8 tonnes of watermelon seed is imported from countries where *A. citrulli* occurs, this would equate to between approximately 24 million and 96 million seeds.

In terms of the number of consignments imported by NL, at a very minimum each cell in Table 2.05a represents at least one consignment. Thus in 2008 there would have been at least 6 consignments of cucurbit seeds from countries where *A. citrulli* occurs (one from each of China, Thailand, USA, Israel, Japan and Taiwan). And in 2010 there was a minimum of 3 consignments (China, USA and Israel). Whilst in 2008 the imports from USA, Taiwan and Israel could well have been single consignments of between 1 kg and 51 kg, the import from China is less likely to have been a single consignment of over 16 tonnes. Equally it is very unlikely that the Chinese imports were 16,000 consignments of just over 1kg each.

Taking the weight of the cucurbit seed imported by NL from Thailand, USA, Israel, Japan and Taiwan between 2008 and 2010, then ranking the data in order gives an ordered list of: 1, 35, 37, 51, 75, 100, 122, 192, 209 and 939kg. Excluding the outliers of 1 and 939, the mean weight of seed from these countries is approximately 102kg. Assuming that the typical seed consignment (lot) of imported cucurbit seed weighs approximately 100kg, and that records of more than 100kg imports represent multiple consignments, then we can estimate for example that approximately 162 consignments of cucurbit seed were imported by NL from China in 2008. Applying such an approach across table 2.05a suggest that between 2008 and 2010 an average of approximately 170 consignments of cucurbit seed were

imported by NL from countries where *A. citrulli* occurs. Recognising that NL is very likely to be the major EU importer, importing the majority of consignments, it can be estimated that the EU as a whole imports perhaps around 200 consignments of cucurbit seed per year from countries where *A. citrulli* occurs.

### Uncertainties regarding the quantity of commodity imported annually

Although in the EU, and generally elsewhere in the world, there is very little statistical information collected specifically about the trade in cucurbit seeds, we can be confident that the magnitude of EU imports is a fraction of total vegetable seed imports of up to 100,000 tonnes per year (Table 2.05a 1) and from countries where *A. citrulli* occurs is a fraction of 44,000 tonnes (Table 2.05a 2), assuming that the EU does source cucurbit seed from such countries.

Specific information from the Netherlands enabled approximations to be made regarding the number of cucurbit seeds imported. However, the proportion of cucurbit seed that is watermelon, and the most likely source of the pathogen, is unknown.

Based on NL data for 2008 – 2010, crude calculations have been used to estimate that approximately 200 consignments of cucurbit seed are imported into the EU each year from countries where *A. citrulli* occurs

### Conclusions

The quantity of watermelon seed imported into EU MS from countries where *A. citrulli* occurs is assumed to be primarily in the order of 10 million to 100 million seeds per year.

The table below uses a log scale for categorising the volume of commodity imported / number of seeds / plants imported into the EU.

<b>2.05a Quantity of annual imports</b> (Examples provided based on a log scale of millions of seeds).			
<b>Rating</b>	<b>Number of seeds imported per year</b>	<b>Justification summary</b>	<b>Probability Assignment <sup>1</sup></b>
<b>Very low</b>	< 1 million	-	
<b>Low</b>	1 to 10 million	There is some uncertainty around the calculations	9%
<b>Medium</b>	10 to 100 million	Assuming 2 to 8 tonnes of susceptible seed is imported, this equates to 24 to 96 million seeds	90%
<b>High</b>	100 to 1,000 million	There is some uncertainty around the calculations	1%
<b>Very high</b>	>1,000 million	-	
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

## 2.06a Likelihood of transfer via pathway

Consider the likelihood that the commodity will be distributed and subsequently allow the pest to transfer to a suitable host. For example, consider the geographic location of likely markets and the proportion of the commodity that is likely to move to locations where the pest could transfer to a host. Even if infested commodities are shipped to areas where environmental factors allow establishment, unless the pest can locate a host, establishment will not be possible. Consider the intended use of the commodity, e.g. plants for planting or produce for processing and consumption; likelihood of transfer from by-products of processing, or disposal of the commodity in the vicinity of suitable hosts; the pest's ability to disperse and whether vectors provide a route from the pathway to a host; the time of year at which import takes place.

If possible consider the likelihood that sufficient numbers of the pest will transfer from the pathway to a suitable host in order to potentially initiate a new population. The reproductive strategy of the pest should be taken into account. Alternatively consider the potential number of "transfer events" that could occur per infested/contaminated consignment that has entered. See table for descriptions.

**Information / evidence:** *Provide reasoning then give judgment*

The commodity is seed for planting so there is a very high likelihood that infected seed would transfer to a growing crop if distributed to the grower.

### Uncertainties regarding likelihood of transfer

None

### Conclusions

If a seedlot is infected there is a very high likelihood of transfer.

2.06a: Likelihood pest will transfer in sufficient numbers to a host			
Rating	Description (likelihood of pest transfer is ....)	Justification summary	Probability Assignment <sup>1</sup>
Very low	< 0.01% (less than one in ten thousand contaminated lots will provide transfer opportunities)		%
Low	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand contaminated lots will provide transfer opportunities)		%
Medium	Between 0.1% - 1% (between one in one thousand and one in one hundred contaminated lots will provide transfer opportunities)		%
High	Between 1% and 10% (between one in one hundred and one in ten contaminated lots will provide transfer opportunities)		%
Very high	> 10% (more than one in ten contaminated lots will provide transfer opportunities)	Pathway is seeds for planting. If contaminated transfer highly likely	100%
	Check sum =		100%

<sup>1</sup> Spread your judgment according to your belief / evidence

### **2.07a Likelihood of entry and transfer via pathway “a” (Seeds of cucurbits, especially watermelon, from countries where *Acidovorax citrulli* occurs)**

Use the BBN to combine the scores to questions 2.01 to 2.06, which all relate to the likelihood of pest entry then transfer. Present it as Figure 2.07.

The result of combining scores to individual questions 2.01(a) to 2.05(a), that relate to likelihood of entry are combined with score for likelihood of transfer 2.06(a) using a BBN to provide an assessment of potential for entry and transfer for the pathway and is shown in Figure 2.07 a. It suggests that the potential for entry and transfer via the pathway “Seeds of cucurbits, especially watermelon, from countries where *Acidovorax citrulli* occurs” has some uncertainty given that the overall likelihood spans three of the five categories, i.e. likelihood of entry and transfer is between “low” and “high”. The most likely likelihood is “medium”. Given the few occurrences where bacterial fruit blotch have been associated with seed in the EU, the model outcome seems appropriate and in line with assessors judgment.

Assuming 200 consignments of potentially infected cucurbit seed are imported into the EU each year (2.05a) and there is an 80% chance that between 0.1 and 1% of them are contaminated (2.01a) and a 20% chance that between 1% and 10% are contaminated, then it would be expected that between 0.2 and 4 consignments of contaminated seed arrive in the EU each year (Box 2.07a).

#### **Box 2.07a: Estimating the number of contaminated cucurbit seed lots / consignments entering the EU**

Total consignments = 200

80% likelihood that 0.1 to 1% are contaminated =  $200 \times 0.8 \times 0.001$  to  $200 \times .08 \times 0.01$   
= 0.16 to 1.6

20% likelihood that 1 to 10% are contaminated =  $200 \times 0.2 \times 0.01$  to  $200 \times .02 \times 0.1$   
= 0.4 to 4.0


#### **Uncertainties regarding likelihood of entry and transfer**

There is a lack of detail regarding the species of cucurbit seed imported into the EU and of its origin. As noted previously the time of year material is moved, volumes, routes and modes of transport used are also unknown. Whether or not seed is tested, and whether *A. citrulli* has been intercepted in the EU on seed previously is unknown. It is assumed not to have been intercepted on seed in the EU. However seed testing is not mandatory. Assumptions had to be made about the environmental conditions in which seed is transported.

#### *Conclusions*

There is an ongoing and unregulated international trade in cucurbit seed. The EU sources such seed from countries where *A. citrulli* occurs. Presumably such trade has been ongoing for many years. There have only been a few incidents of bacterial fruit blotch outbreaks in the EU and none have been proven to link to seed imports. The BBN model (described elsewhere) suggests that the likelihood of entry and transfer via seed is most likely considered “medium”. The risk elements (questions) that have most influence on the final outcome of the model pathway are the likelihood of association at origin (Q 2.01a) and the quantity imported (Q 2.05a). If more information were available to help reduce uncertainty surrounding these questions, then the uncertainty involved in this pathway would be reduced.

**Figure 2.07a:** Model output from BBN indicating likelihood of entry of *Acidovorax citrulli* via seeds of cucurbits, especially watermelon, from countries where *Acidovorax citrulli* occurs.



**Pathway 2: Seedlings of cucurbits** (especially watermelon, from countries where *Acidovorax citrulli* occurs).

**2.01b Likelihood of the pest being associated, spatially and temporally, with the pathway at origin**

(There must be some likelihood of association otherwise there is no pathway).

Take into account pre-harvest cultivation and husbandry practices such as existing pest management measures, choice of cultivar, and applications of plant protection products. If phytosanitary measures (i.e. statutory risk reduction measures) are already in place against this or other pests at origin, specify whether these are being taken into account or not.

**Information / evidence:** *Provide reasoning then give judgment*

The EU does not provide data on exports and imports of cucurbit seedlings (plants for planting) hence there is very little data available about the EU import and trade of cucurbit transplants. However, anticipating the lack of statistics, the Prima phacie case study consortium developed a questionnaire for EU MS to address some anticipated information gaps (Annex 1). Part of the questionnaire was designed to determine which EU MS import seedlings and from where such transplants are obtained.

Only three of the 13 EU MS that replied to the questionnaires stated that cucurbit seedlings had indeed been imported. The Czech Republic had sourced cucurbit seedlings from China, a country where *A. citrulli* is known to occur. Italy has imported seedlings but did not know their origin and Germany had received seedlings from the Netherlands. Although the later movement of seedlings from the Netherlands to Germany is not strictly an “import”, since it was within the single market, it is possible that the seedlings had come from a Third Country via the Netherlands.

The pathogen has been found in association with seedlings of cucurbits in trade; the first report of what is now called *Acidovorax citrulli* was in 1965, in Georgia, USA when an unidentified seedborne bacterium was isolated from necrotic watermelon cotyledons originating in Turkey (Webb & Goth, 1965; Bahar & Burdman, 2010). Hungary had an outbreak of *A. citrulli* in July 2007. The source of infection was unclear, but it was noted that grafted watermelon transplants had been imported from Turkey (Palkovics *et al.*, 2008). Latin & Hopkins (1995) reported that the outbreak in Iowa state was linked to purchase of transplants from an Indiana grower who suffered outbreak of the disease in his transplant facilities and in his fields.

The source of infection for most seedlings is the seed itself. It was shown by Shirakawa *et al.* (2003) that under high humidity infested seeds with only 1cfu of *A. citrulli* have a high probability of resulting in diseased germinated seedlings. Glasshouse conditions, where seedlings are likely to be grown include high temperatures and high relative humidity, which alongside high populations of plants is conducive to seedling transmission and rapid disease spread (Walcott, 2005). As few as 1-2 seedlings can lead to widespread disease (Schaad, 2003).

**Uncertainties regarding likelihood of the pest being associated with the pathway at origin**

Data on the specific area of origin of cucurbit seedlings is unavailable, so it is unknown how easy it is to source plant material which can be guaranteed free of infection at origin. Level of association is unclear.

## Conclusions

There is a likelihood of association at origin, but level is unclear.

<b>2.01b: Likelihood of association with the pathway at origin</b>			
<b>Rating</b>	<b>Description</b> (likelihood of association is ...)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the commodity are likely to be contaminated / infested)	Level of infected seedlings in trade is unknown – but there have been occurrences in trade.  Lack of outbreaks widely across the EU suggests that association is not likely high or very high	-
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> are likely to be contaminated / infested)		25%
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the commodity are likely to be contaminated / infested)		70%
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the commodity are likely to be contaminated / infested)		5%
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the commodity are likely to be contaminated / infested)		-
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence.

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

*The assessment of pest entry and transfer (2.01 -2.06) is based on considering "lots".  
However, if an alternative unit would be more appropriate then describe the chosen unit and ordinal scale using five categories.*

## **2.02b Likelihood of surviving postharvest treatments / measures (before pest entry into risk assessment area)**

*Given that a proportion of lots/ consignments may be infested / contaminated, consider the proportion of infested/contaminated lots that are likely to remain infested/contaminated after any manipulation, handling or specific phytosanitary treatment to which the commodity is subjected. Examples of postharvest treatments include culling, washing, chemical treatment and cold storage. If there are no post harvest treatments the likelihood of survival should probably be considered “very high”.*

*If post-harvest phytosanitary measures (i.e. statutory risk reduction measures) are already in place, specify whether these are being taken into account or not.*

**Information / evidence:** *Provide reasoning then give judgment*

*Acidovorax citrulli* is not specifically mentioned in the EC Marketing Directive 92/33/EEC of 28<sup>th</sup> April 1992, although there is a requirement that vegetable propagating and planting material (other than seed) of *Cucumis melo* (melon), *Cucumis sativus* (gerkin), *Cucurbita pepo* (courgette), *Cucurbita maxima* (gourd) be substantially pest free. In addition, there are no specific requirements for this pathogen in the EU Plant Health Legislation.

Seedlings may show symptoms of infection, especially if grown in warm conditions of high humidity. Obviously symptomatic plants are likely to be rejected before being sent to customers, however, not all infected seedlings will show symptoms.

### **Uncertainties regarding likelihood of the pest surviving postharvest treatments/ measures (before pest entry into the risk assessment area)**

The level of seedlings in which infection may remain latent is unknown.

## Conclusions

Lower likelihood of association than with seed as some infected seedlings are likely to be weeded out, but symptoms may not show in all cases (latent infections) and their expression may be affected by the growing conditions.

<b>2.02b: Likelihood that an infested/contaminated commodity remains infested/contaminated after existing postharvest treatments/measures</b>			
<b>Rating</b>	<b>Description</b> (likelihood of remaining infested/contaminated / pest survival is ....)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Some infected seedlings likely to be weeded out, but not all will show symptoms and hence remain on the pathway.	45%
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		45%
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		10%
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

### 2.03b Likelihood of surviving storage and transport

Given that a proportion of lots/ consignments may still be infested / contaminated, estimate the proportion of lots that are likely to remain infested/contaminated because the pest can survive storage and transport; consider speed and conditions of transport and duration of the life cycle of the pest in relation to time in storage and transport, commercial procedures (e.g. refrigeration) applied to consignments in the country of origin, during shipping, and in the country of destination, that could affect the likelihood of pest survival. Take into account previous live interceptions on this or similar pathways (see 1.6).

If phytosanitary measures (i.e. statutory risk reduction measures) are already in place which act on the likelihood of pest survival during storage and transport, specify whether these are being taken into account or not.

#### Information / evidence: *Provide reasoning then give judgment*

There is no data available on the length of time taken for transport and storage of this commodity. However it is likely to be short given the fragile nature of young plants and the need for them to be planted and grown on. No known reason why pest could not survive.

Using artificial inoculations of melon plants, Silva *et al.* (2006), reported an epiphytial survival of *A. citrulli* on leaves of 54 days, in greenhouse as well as in the field. The epiphytic population increased initially and decreased after a certain period of time. In a glasshouse, the bacterial population on the roots and rhizosphere decreased with time.

#### Uncertainties regarding likelihood of the pest surviving storage and transport

None

#### Conclusions

Young plants are likely to be shipped and stored for only a short time. *A. citrulli* highly likely to survive.

2.03b: Likelihood of surviving storage and transport			
Rating	Description (likelihood of remaining contaminated / pest survival is ....)	Justification summary	Probability Assignment <sup>1</sup>
Very Low	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
Low	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
Medium	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
High	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
Very High	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Storage and shipping time likely to be short	100%
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup>Lot: a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

#### **2.04b Likelihood of pest surviving current phytosanitary procedures at the point of entry or elsewhere in the risk assessment area**

*Given that a proportion of lots may still be infested / contaminated, estimate the proportion of lots that are likely to remain infested/contaminated because the pest survives existing phytosanitary procedures e.g. it is not detected at entry and/ or it can survive any existing phytosanitary procedures within the pest risk assessment area. Take into account the intensity of sampling and inspection and ease of detecting and distinguishing the pest from other organisms.*

#### **Information / evidence:** *Provide reasoning then give judgment*

Within EU phytosanitary legislation, cucurbit seedlings are considered as “plants for planting” and if they originate in Third Countries (i.e. non EU MS) they need a phytosanitary certificate to indicate that the seedlings are free from EU listed quarantine pests.

Pathogen free transplants can be produced through a thorough inspection of the plant material before introduction to the growing area (glasshouse or field). However, since *A. citrulli* is not a listed quarantine pest (i.e. not a listed “harmful organism” in the Plant Health Directive) cucurbit seedlings contaminated with *A. citrulli* could still enter the EU with a valid phytosanitary certificate.

Some cucurbit plants of planting are further regulated by 2000/29 EC, i.e. (i) *Cucumis* spp. (e.g. melons, cucumbers and gherkins) are listed in Annex IV of the Plant Health Directive (2000/29 EC, Annex IV, 32.1) with regard to leaf mining pests (*Amauromyza* and *Liriomyza* spp.). Phytosanitary inspectors examining imported *Cucumis* seedlings would therefore focus on looking for leaf damage and could perhaps see symptomatic lesions caused by *A. citrulli*. (ii) melon plants (*Cucumis melo*) are listed in annex II of the Plant Health Directive (2000/29/EC, Annex II. part A, chapter II, paragraph d, point 15) with regard to a viral pathogen (*Tomato Spotted Wilt Virus*). However, rather than *Cucumis*, it is *Citrullus lanatus* (watermelon) plants for planting that have been implicated in previous outbreaks of bacterial fruit blotch. For example watermelon seedlings from Turkey were linked with the original outbreak in Georgia in the 1960s and the more recent outbreak in Hungary also followed the import of watermelons seedlings from Turkey (Palkovics, 2008). However, there are no special requirements within the Plant Health Directive regarding watermelon plants for planting.

#### **Uncertainties regarding likelihood of the pest surviving current phytosanitary procedures**

Although general phytosanitary checks would be conducted on seedling samples at import, it could be difficult to detect an infected consignment at low levels of infestation.

### Conclusions

Only basic phytosanitary procedures are currently required (visual inspection), therefore continued survival is very likely.

<b>2.04b: Likelihood of pest surviving current phytosanitary procedures</b>			
<b>Rating</b>	<b>Description</b> (likelihood of remaining contaminated/ pest survival is ....)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		%
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Checks possible, but not required	100%
	Check sum =		100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

## 2.05b Quantity of commodity imported annually

*Quantity of commodity imported annually: The likelihood that a pest will enter depends on the amount of the potentially-infested commodity that is imported. For qualitative pest risk assessments, the amount of commodity imported is estimated in units of tonnes, or other metric such as standard 40 foot long shipping containers.*

*If the quantity of commodity imported is better described using alternative units, such as the number of plants for planting, assessors should devise a 5 level scale and provide some reasoning to support use of the scale.*

### **Information / evidence:** *Provide reasoning then give judgment*

Prima phacie case study consortium developed a questionnaire for EU MS to address some anticipated information gaps (Annex 1). Part of the questionnaire was designed to determine which EU MS import seedlings and from where such transplants are obtained. There are no figures for quantities, but very few countries responded that seedlings were imported and quantities are likely to be less than for seed. Countries are more likely to grow their own plants from seed.

The Netherlands have imported cucurbit seedlings from Israel (Dirk Jan van der Gaag, pers. comm.).

### **Uncertainties regarding the quantity of commodity imported annually**

No data is available to support import assumptions.

### *Conclusions*

The quantity of seedlings import is assumed to be less than that of seed, and is assumed mainly “low” as indicated in table 2.05 b below. We assume most EU countries grow cucurbits using EU or imported seed. There is considerable uncertainty around the number of plants / seedlings imported. However, use of a log scale facilitates uncertainty being accounted for.

<b>2.05b Quantity of annual imports</b> (Examples provided based on a log scale of millions of plants / seedlings).			
<b>Rating</b>	<b>Number of plants / seedlings imported per year</b>	<b>Justification summary</b>	<b>Probability Assignment <sup>1</sup></b>
<b>Very low</b>	< 1 million		9%
<b>Low</b>	1 to 10 million		90%
<b>Medium</b>	10 to 100 million		1%
<b>High</b>	100 to 1,000 million		-
<b>Very high</b>	>1,000 million		-
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

**2.06b Likelihood of transfer via the pathway of cucurbit seedlings, especially watermelon, from countries where *Acidovorax citrulli* occurs.**

Consider the likelihood that the commodity will be distributed and subsequently allow the pest to transfer to a suitable host. For example, consider the geographic location of likely markets and the proportion of the commodity that is likely to move to locations where the pest could transfer to a host. Even if infested commodities are shipped to areas where environmental factors allow establishment, unless the pest can locate a host, establishment will not be possible. Consider the intended use of the commodity, e.g. plants for planting or produce for processing and consumption; likelihood of transfer from by-products of processing, or disposal of the commodity in the vicinity of suitable hosts; the pests ability to disperse and whether vectors provide a route from the pathway to a host; the time of year at which import takes place.

If possible consider the likelihood that sufficient numbers of the pest will transfer from the pathway to a suitable host in order to potentially initiate a new population. The reproductive strategy of the pest should be taken into account. Alternatively consider the potential number of “transfer events” that could occur per infested/contaminated consignment that has entered. See table for descriptions.

**Information / evidence:** *Provide reasoning then give judgment*

The commodity is plants for planting, therefore transfer to growing host is highly likely.

**Uncertainties regarding likelihood of transfer**

None.

*Conclusions*

No restrictions on transfer to host.

<b>2.06b: Likelihood pest will transfer in sufficient numbers to a host</b>			
<b>Rating</b>	<b>Description (likelihood of pest transfer is ....)</b>	<b>Justification summary</b>	<b>Probability Assignment <sup>1</sup></b>
<b>Very low</b>	< 0.01% (less than one in ten thousand contaminated lots will provide transfer opportunities)		%
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand contaminated lots will provide transfer opportunities)		%
<b>Medium</b>	Between 0.1% - 1% (between one in one thousand and one in one hundred contaminated lots will provide transfer opportunities)		%
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten contaminated lots will provide transfer opportunities)		%
<b>Very high</b>	> 10% (more than one in ten contaminated lots will provide transfer opportunities)	Infected seedlings are hosts themselves.	100%
	Check sum =		100%

<sup>1</sup> Spread your judgment according to your belief / evidence

**2.07b Likelihood of entry and transfer via pathway “b”, cucurbit seedlings, especially watermelon, from countries where *Acidovorax citrulli* occurs.**

*Use the BBN to combine the scores to questions 2.01 to 2.06, which all relate to the likelihood of pest entry then transfer. Present it as Figure 2.07.*

The result of combining scores to individual questions 2.01(b) to 2.05(b), that relate to likelihood of entry is combined with score for likelihood of transfer 2.06(b) using a BBN to provide an assessment of potential for entry and transfer for the pathway and is shown in Figure 2.07 b. It suggests that the potential for entry and transfer via cucurbit seedlings, especially watermelon, from countries where *Acidovorax citrulli* occurs is most likely “low”.

**Uncertainties regarding likelihood of entry and transfer**

There is a lack of detail regarding the species of cucurbit seedlings imported into the EU and of its origin. As noted previously the time of year material is moved, volumes, routes and modes of transport used are also unknown.

Whether or not seedlings are tested, and whether *A. citrulli* has been intercepted in the EU on seedlings is unknown. It is assumed not to have been intercepted on seedlings.


*Conclusions*

There is some regulated international trade in cucurbit seedlings (plants for planting) although EU legislation focuses on *Cucumis* species rather than *Citrullus* which is more likely to carry the pathogen. Nevertheless the EU sources *Citrullus* from countries where *A. citrulli* occurs. Presumably such trade has been ongoing for many years. There have been a few outbreaks of bacterial fruit blotch in Europe where the import of seedlings from a Third Country was implicated. However, no clear link has been established.

The BBN model (described elsewhere) suggests that the likelihood of entry and transfer via seedlings ranges between “very low” and “medium”, and is most likely “low”. Based on a presumed history of trade since at least the 1960s when seedlings from Turkey were implicated with an outbreak in Georgia, and a lack of interception evidence (recognising that watermelon seedlings are not regulated) and the few EU outbreaks, the model output is broadly in line with the view of assessors.

*Nevertheless the risk elements considered in test Method 4b, and how they operate within the BBN would benefit from further study.*

**Figure 2.07b:** Model output from BBN indicating likelihood of entry of *Acidovorax citrulli* via seedlings of cucurbits, especially watermelon, from countries where *Acidovorax citrulli* occurs.



**Pathway 3: Fruit of cucurbits (especially watermelon, imported from countries where *Acidovorax citrulli* occurs)**

**2.01c Likelihood of the pest being associated, spatially and temporally, with the pathway at origin**

(There must be some likelihood of association otherwise there is no pathway).

Take into account pre-harvest cultivation and husbandry practices such as existing pest management measures, choice of cultivar, and applications of plant protection products. If phytosanitary measures (i.e. statutory risk reduction measures) are already in place against this or other pests at origin, specify whether these are being taken into account or not.

**Information / evidence:** *Provide reasoning then give judgment*

The EU imports a variety of cucurbit fruit from many parts of the world. An examination of the combined nomenclature (CN) codes used by custom authorities reveals that imports of cucurbit fruit are categorised into a variety of codes, three of the groupings were used to assess imports of fresh watermelons (CN 08071010), fresh cantaloupe and other melons (CN 08071090) and fresh or chilled cucumbers & gherkins (CN07070005).

Tables 2.01c a to d show the countries that have been the biggest suppliers of cucurbit fruit to the EU 27 over the past 3 years (2008 – 2010). Those countries where *A. citrulli* is present are highlighted. In all cases the EU has sourced cucurbit fruit from countries where the pathogen occurs e.g. watermelon from Brazil, Costa Rica and Turkey.

<b>Table 2.01c a: EU 27 imports of watermelons from non EU countries 2008 – 2010 (tonnes)</b>						
Source of import \ Year	2008	2009	2010	3 year mean	% of total	cumulative %
Panama	32,526	24,156	33,811	30,164	16.2	16.2
Brazil *	32,131	30,957	23,769	28,953	15.5	31.7
Costa Rica *	23,567	28,185	30,957	27,570	14.8	46.5
Macedonia	29,218	17,316	16,810	21,115	11.3	57.9
Turkey *	32,857	4,241	13,694	16,931	9.1	67.0
Ukraine	3,917	19,096	25,197	16,070	8.6	75.6
Tunisia	17,264	9,768	14,246	13,759	7.4	83.0
Russia	10,899	10,027	6,883	9,269	5.0	88.0
Morocco	5,640	7,646	3,369	5,552	3.0	90.9
Jordan	4,269	5,172	5,525	4,988	2.7	93.6
Egypt	3,743	4,106	5,146	4,332	2.3	95.9
Peru	904	2,280	2,975	2,053	1.1	97.0
Iran *	767	1,632	1,377	1,259	0.7	97.7
28 other non EU countries	3,885	2,643	6,188	4,239	2.3	100.0
sum	201,587	167,223	189,946	186,252	100.0	

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

**Table 2.01c b:** EU 27 imports of melons (excluding watermelons) from non EU countries 2008 – 2010 (tonnes)

Year	2008	2009	2010	3 year mean	% of total	cumulative %
Brazil *	193,025	173,423	168,903	178,450	53.2	53.2
Morocco	56,544	55,310	54,316	55,390	16.5	69.7
Costa Rica *	50,969	43,176	59,656	51,267	15.3	84.9
Honduras	17,380	23,171	21,471	20,674	6.2	91.1
Panama	20,128	13,222	11,268	14,873	4.4	95.5
Israel *	4,073	4,606	4,630	4,436	1.3	96.8
46 other non EU countries	9,943	10,280	11,621	10,615	3.2	100.0
sum	352,062	323,187	331,866	335,705	100.0	

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

**Table 2.01c c:** EU 27 imports of cucumbers (& gherkins) from non EU countries 2008 – 2010 (tonnes)

Year	2008	2009	2010	3 year mean	% of total	cumulative %
Turkey *	17,648	20,112	20,658	19,473	66.3	66.3
Macedonia	3,351	2,734	3,890	3,325	12.6	78.9
Morocco	2,986	3,249	2,350	2,862	11.2	90.1
Jordan	1,250	2,377	3,475	2,367	4.7	94.8
Bosnia & Herzegovina	628	1,068	485	727	2.4	97.2
Syria	56	739	573	456	0.2	97.4
Egypt	299	310	264	291	1.1	98.5
Israel *	246	232	157	212	0.9	99.4
49 other non EU countries	150	703	819	557	0.6	100.0
sum	26,612	31,525	32,670	30,269	100.0	

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

**Table 2.01c d:** EU 27 imports of cucurbit fruit from Third countries where *Acidovorax citrulli* occurs (mean quantity imported 2008 – 2010, extracted from Tables 2.05c a to c) (tonnes)

Commodity	Potential source	Mean quantity imported 2008 - 2010
Watermelons	Brazil	28,953
	Costa Rica	27,570
	Turkey	16,931
	Sub total	73,454
Melons (excluding watermelons)	Brazil	178,450
	Costa Rica	51,267
	Israel	4,436
	Sub total	234,153
Cucumbers & gherkins	Turkey	19,473
	Israel	212
	Sub total	19,685
	Sum	327,292

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

Generally, evidence of association of a pathogen with host commodities can be seen in interception reports. However, because *A. citrulli* is not currently regulated in the EU, there is no formal requirement to share information on any such interceptions. Nevertheless, there is one report of Wall *et al.* (1990) where it is reported that fruits were intercepted by quarantine persons inspecting a watermelon shipment from Tinian to Guam.

**Uncertainties regarding likelihood of the pest being associated with the pathway at origin**

Level of infection in the fruit in the exporting countries is unknown.

*Conclusions*

A high level of cucurbit fruit is imported to the EU from countries where the pest is present, but the proportion of this that is likely to be infected will be considerably lower

<b>2.01c: Likelihood of association with the pathway at origin</b>			
<b>Rating</b>	<b>Description (likelihood of association is ....)</b>	<b>Justification summary</b>	<b>Probability Assignment <sup>1</sup></b>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the commodity are likely to be contaminated / infested)		10%
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> are likely to be contaminated / infested)		80%
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the commodity are likely to be contaminated / infested)		10%
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the commodity are likely to be contaminated / infested)		-
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the commodity are likely to be contaminated / infested)		-
Check sum =			100%

<sup>1</sup> Spread your judgment according to your belief / evidence.

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

*The assessment of pest entry and transfer (2.01 -2.06) is based on considering "lots".  
However, if an alternative unit would be more appropriate then describe the chosen unit and ordinal scale using five categories.*

**2.02c Likelihood of surviving postharvest treatments / measures (before pest entry into risk assessment area)**

Given that a proportion of lots/ consignments may be infested / contaminated, consider the proportion of infested/contaminated lots that are likely to remain infested/contaminated after any manipulation, handling or specific phytosanitary treatment to which the commodity is subjected. Examples of postharvest treatments include culling, washing, chemical treatment and cold storage. If there are no post harvest treatments the likelihood of survival should probably be considered “very high”.

If post-harvest phytosanitary measures (i.e. statutory risk reduction measures) are already in place, specify whether these are being taken into account or not.

**Information / evidence:** *Provide reasoning then give judgment*

Any lot of diseased fruit is likely to be unmarketable and will not be traded. However, fruit can be infected without visible symptoms (Walcott *et al.*, 2003; Lessl *et al.*, 2007; Bahar *et al.*, 2009). Other than sorting and grading fruit in terms of quality, no specific measures are in place to reduce the likelihood that contaminated fruit will be removed from the pathway and the proportion of asymptomatic infested fruit is likely to remain quite high.

**Uncertainties regarding likelihood of the pest surviving postharvest treatments/ measures (before pest entry into the risk assessment area)**

The proportion of fruit that would be infected, but without visible symptoms is unknown.

*Conclusions*

Obviously diseased fruit are unmarketable and will be excluded. A proportion will be infected and not show visible symptoms.

<b>2.02c: Likelihood that an infested/contaminated commodity remains infested/contaminated after existing postharvest treatments/measures</b>			
<b>Rating</b>	<b>Description</b> (likelihood of remaining infested/contaminated / pest survival is ....)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		-
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Assume that there is a good chance that 1 to 10% of contaminated fruit do not show symptoms	50%
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Assume that there is a good chance that more than 10% of contaminated fruit do not show symptoms	50%
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

### 2.03c Likelihood of surviving storage and transport

Given that a proportion of lots/ consignments may still be infested / contaminated, estimate the proportion of lots that are likely to remain infested/contaminated because the pest can survive storage and transport; consider speed and conditions of transport and duration of the life cycle of the pest in relation to time in storage and transport, commercial procedures (e.g. refrigeration) applied to consignments in the country of origin, during shipping, and in the country of destination, that could affect the likelihood of pest survival. Take into account previous live interceptions on this or similar pathways (see 1.6).

If phytosanitary measures (i.e. statutory risk reduction measures) are already in place which act on the likelihood of pest survival during storage and transport, specify whether these are being taken into account or not.

#### Information / evidence: *Provide reasoning then give judgment*

Survival of the pest during transport / storage depends on the transport / storage time and conditions, but transport and storage of such perishable commodities is likely to be relatively short and conditions cool and dry. The pathogen will continue to survive in such conditions.

#### Uncertainties regarding likelihood of the pest surviving storage and transport

None.

#### Conclusions

Survival is very likely.

2.03c: Likelihood of surviving storage and transport			
Rating	Description (likelihood of remaining contaminated / pest survival is ....)	Justification summary	Probability Assignment <sup>1</sup>
Very Low	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
Low	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
Medium	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
High	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
Very High	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	No reason for the pathogen not to survive transport or storage.	100%
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup>Lot: a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

**2.04c Likelihood of pest surviving current phytosanitary procedures at the point of entry or elsewhere in the risk assessment area**

*Given that a proportion of lots may still be infested / contaminated, estimate the proportion of lots that are likely to remain infested/contaminated because the pest survives existing phytosanitary procedures e.g. it is not detected at entry and/ or it can survive any existing phytosanitary procedures within the pest risk assessment area. Take into account the intensity of sampling and inspection and ease of detecting and distinguishing the pest from other organisms.*

**Information / evidence:** *Provide reasoning then give judgment*

Cucurbit fruit are not regulated by the EC plant health directive and so no specific phytosanitary measures are in place. *A. citrulli* is not a listed quarantine pest and there are no current phytosanitary requirements in place to inhibit its introduction on cucurbit fruit.

**Uncertainties regarding likelihood of the pest surviving current phytosanitary procedures**

None

*Conclusions*

<b>2.04c: Likelihood of pest surviving current phytosanitary procedures</b>			
<b>Rating</b>	<b>Description</b> (likelihood of remaining contaminated/ pest survival is ....)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very Low</b>	< 0.01% (less than one in ten thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
<b>Medium</b>	Between 0.1% and 1% (between one in one thousand and one in one hundred lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)		
<b>Very High</b>	> 10% (more than one in ten lots <sup>2</sup> of the original commodity are likely to remain contaminated / infested)	Without measures in place there is no reason for pathogen not to survive.	100%
Check sum =			100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<sup>2</sup>**Lot:** a number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (ISPM no. 5, IPPC, 2007).  
A consignment may be several lots or a single lot.

### 2.05c Quantity of commodity imported annually

Quantity of commodity imported annually: The likelihood that a pest will enter depends on the amount of the potentially-infested commodity that is imported. For qualitative pest risk assessments, the amount of commodity imported is estimated in units of tonnes, or other metric such as standard 40 foot long shipping containers.

If the quantity of commodity imported is better described using alternative units, such as the number of plants for planting, assessors should devise a 5 level scale and provide some reasoning to support use of the scale.

#### Information / evidence: Provide reasoning then give judgment

The EU imports a variety of cucurbit fruit from many parts of the world. Tables 2.01c a to d (see previous) show the countries that have been the biggest suppliers of cucurbit fruit to the EU 27 over the past 3 years (2008 – 2010). In the 3 years 2008 – 2010 EU MS have sourced just over 40% of watermelon fruit from countries where *A. citrulli* occurs (Table 2.05c a). Over the same period almost 70% of other imported melons have come from Brazil, Costa Rica and Israel (Table 2.05c b), where *A. citrulli* also occurs.

Table 2.05c a: Mean EU 27 imports of watermelons from countries where <i>A. citrulli</i> is present (marked*) and other sources ( <i>A. citrulli</i> assumed absent) 2008 – 2010 (tonnes)		
Source of import	Mean annual imports (tonnes)	% of total
Brazil *	28,953	
Costa Rica *	27,570	
Turkey *	16,931	
Iran *	1,259	
Sub total	74,713	40.1
Panama	30,164	
Macedonia	21,115	
Ukraine	16,070	
Tunisia	13,759	
Russia	9,269	
Morocco	5,552	
Jordan	4,988	
Egypt	4,332	
Peru	2,053	
28 other non EU countries	4,239	
Sub total	111,541	59.9
sum	186,252	100.0

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

<b>Table 2.05c b: Mean EU 27 imports of melons (excluding watermelons) from countries where <i>A. citrulli</i> is present (marked*) and other sources (<i>A. citrulli</i> assumed absent) 2008 – 2010 (tonnes)</b>		
Source of import	Mean annual imports (tonnes)	% of total
Brazil *	178,450	
Costa Rica *	51,267	
Israel *	4,436	
Sub total	234,153	69.7
Morocco	55,390	
Honduras	20,674	
Panama	14,873	
46 other non EU countries	10,615	
Sub total	101,552	30.3
sum	335,705	100.0

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

<b>Table 2.05c c: Mean EU 27 imports of cucumbers &amp; gherkins from countries where <i>A. citrulli</i> is present (marked*) and other sources (<i>A. citrulli</i> assumed absent) 2008 – 2010 (tonnes)</b>		
Source of import	Mean annual imports (tonnes)	% of total
Turkey *	19,473	
Israel *	212	
Sub total	19,685	65.0
Macedonia	3,325	
Morocco	2,862	
Jordan	2,367	
Bosnia & Herzegovina	727	
Syria	456	
Egypt	291	
49 other non EU countries	557	
Sub total	10,585	35.0
sum	30,270	

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

<b>Table 2.05c d:</b> EU 27 imports of cucurbit fruit from Third countries where <i>Acidovorax citrulli</i> occurs (mean quantity imported 2008 – 2010, extracted from Tables 2.05c a to c) (tonnes)			
Commodity	Potential source	Mean quantity imported 2008 - 2010	
Watermelons	Brazil	28,953	
	Costa Rica	27,570	
	Turkey	16,931	
	Sub total		73,454
Melons (excluding watermelons)	Brazil	178,450	
	Costa Rica	51,267	
	Israel	4,436	
	Sub total		234,153
Cucumbers & gherkins	Turkey	19,473	
	Israel	212	
	Sub total		19,685
	Sum		327,292

Source: EU Eurostat Easy Comext EU 27 Trade by CN8

### Uncertainties regarding the quantity of commodity imported annually

Unlike the previous pathways (seed and seedlings) there is relatively good data showing that EU MS import cucurbit fruit from countries where *A. citrulli* occurs.

### Conclusions

Over recent years (2008 – 2010) in the order of over 300,000 tonnes of watermelons, melons, cucumbers and gherkins have been imported annually from countries where *A. citrulli* occurs (primarily Brazil, Costa Rica, Turkey and Israel (Table 2.05c, a - c)).

<b>2.05c Quantity of annual imports</b> (Examples provided for tonnes and containers, other units can be used) (If an alternative scale is used, describe each category in the scale)				
Rating	Tonnes imported into PRA area (per year)	Number of containers (per year)	Justification summary	Probability Assignment <sup>1</sup>
Very low	< 100	<10 containers		-
Low	100 – 1,000	10- 100 containers		-
Medium	1,000 -100,000	100 – 10,000 containers		-
High	100,000 – 1,000,000	10,000 – 100,000 containers	Refer to Table 2.05c d	100%
Very high	> 1,000,000	> 100,000 containers		-
			Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

## 2.06c Likelihood of transfer via pathway

Consider the likelihood that the commodity will be distributed and subsequently allow the pest to transfer to a suitable host. For example, consider the geographic location of likely markets and the proportion of the commodity that is likely to move to locations where the pest could transfer to a host. Even if infested commodities are shipped to areas where environmental factors allow establishment, unless the pest can locate a host, establishment will not be possible. Consider the intended use of the commodity, e.g. plants for planting or produce for processing and consumption; likelihood of transfer from by-products of processing, or disposal of the commodity in the vicinity of suitable hosts; the pest's ability to disperse and whether vectors provide a route from the pathway to a host; the time of year at which import takes place.

If possible consider the likelihood that sufficient numbers of the pest will transfer from the pathway to a suitable host in order to potentially initiate a new population. The reproductive strategy of the pest should be taken into account. Alternatively consider the potential number of "transfer events" that could occur per infested/contaminated consignment that has entered. See table for descriptions.

**Information / evidence:** *Provide reasoning then give judgment*

The intended use of fruit is consumption. There is some transfer potential if seeds of diseased fruits were used for planting, but this is highly unlikely to happen, and certainly not on a commercial scale. Transfer of the pathogen from infected fruit to a growing crop seems unlikely unless the fruit is disposed of carelessly, very close to a site of production.

### Uncertainties regarding likelihood of transfer

Whether contaminated fruit is disposed of close to cucurbit production and whether seed from infected fruit is used in cucurbit production.

### Conclusions

Very low likelihood of transfer to a host.

<b>2.06c: Likelihood pest will transfer in sufficient numbers to a host</b>			
<b>Rating</b>	<b>Description (likelihood of pest transfer is ....)</b>	<b>Justification summary</b>	<b>Probability Assignment <sup>1</sup></b>
<b>Very low</b>	< 0.01% (less than one in ten thousand contaminated lots will provide transfer opportunities)	Very unlikely that fruit provides transfer opportunity	100%
<b>Low</b>	Between 0.01% and 0.1% (between one in ten thousand and one in one thousand contaminated lots will provide transfer opportunities)		-
<b>Medium</b>	Between 0.1% - 1% (between one in one thousand and one in one hundred contaminated lots will provide transfer opportunities)		-
<b>High</b>	Between 1% and 10% (between one in one hundred and one in ten contaminated lots will provide transfer opportunities)		-
<b>Very high</b>	> 10% (more than one in ten contaminated lots will provide transfer opportunities)		-
Check sum =			100%

<sup>1</sup> Spread your judgment according to your belief / evidence

### **2.07c Likelihood of entry and transfer via pathway “c” Cucurbit fruit, especially watermelon, imported from countries where *Acidovorax citrulli* occurs.**

*Use the BBN to combine the scores to questions 2.01 to 2.06, which all relate to the likelihood of pest entry then transfer. Present it as Figure 2.07c.*

The result of combining scores to individual questions 2.01(c) to 2.05(c), that relate to likelihood of entry is combined with score for likelihood of transfer 2.06(c) using a BBN to provide an assessment of potential for entry and transfer for the pathway and is shown in Figure 2.07c. It suggests that the potential for entry and transfer via cucurbit fruit, especially watermelon, imported from countries where *Acidovorax citrulli* occurs is “very low”.

#### **Uncertainties regarding likelihood of entry and transfer**

*(Summarise uncertainties from 2.01 to 2.06)*


As is usually the case when assessing any pathway, there is a lack of detailed information. Procedures for harvesting, sorting and grading are unknown. These are steps that could potentially filter out contaminated fruit. Alternatively such procedures could potentially disseminate the pathogen more widely within a consignment through mechanical transmission. How fruit is distributed within the EU is unknown, although the final destination is presumably linked to population density. Disposal of any contaminated fruit is assumed not likely to lead to the pathogen transferring to a site of cucurbit production.

#### *Conclusions*

The EU imports large amounts of cucurbit fruit such as watermelons, melons and cucumbers from countries where *A. citrulli* occurs. However, if any contaminated fruit were to enter the EU, its intended use (consumption and disposal of waste) is not likely to provide opportunities for transfer of the pathogen to sites of cucurbit production.

The results from the BBN model (described elsewhere) as shown in Figure 2.07c suggests that the likelihood of entry and transfer via cucurbit fruit is “very low” and is in line with assessors judgement.

**Figure 2.07c:** Graphical representation of combining scores for questions 2.01c to 2.06c, using a BBN to give an overall likelihood of entry and transfer on pathway c, cucurbit fruit from countries where *A. citrulli* occurs.




If there are multiple pathways, repeat steps 2.01 to 2.07 for each pathway.

There are no more pathways to consider.

## 2.08 Overall likelihood of entry and transfer via all pathways assessed

The likelihood of entry and transfer from all pathways is combined in the BBN, results of which are shown in Figure 2.08 below.

**Figure 2.08:** Overall likelihood of entry and transfer via all pathways



## Conclusions

Figure 2.08 shows the BBN model output from combining scores of entry and transfer for the three pathways assessed, cucurbit seed (2.07a), cucurbit seedlings (2.07b) and cucurbit fruit (2.07c). The range of likelihoods spreads over four of the possible five categories, indicating there is considerable uncertainty. Much uncertainty surrounds the two primary pathways most likely to facilitate entry of *A. citrulli* into the EU so it is not surprising that when all pathways are considered together there is also much uncertainty. Focussing on the most likely of the four categories shown in the bar chart above (2.08) overall likelihood of entry and transfer is considered “medium” to “high”.

Given the history of trade in cucurbit seed, seedlings and fruit from countries where *A. citrulli* occurs and the low number of interceptions reported from around the world, together with the few incidents of outbreaks in the EU, the model output suggests a higher likelihood of entry and transfer than expected. Further study of how pathways are combined in the BBN model is justified.

### **3.00 Potential for pest establishment<sup>6</sup> and extent of spread given entry and transfer**

*Having transferred to a host we next consider whether the pest will survive and reproduce to initiate a population that will establish.*

#### **3.01 Environmental suitability** (particularly considering climate and hosts)

*When introduced to new areas, pests can be expected to behave as they do in their native areas if host plants and climates are similar. Ecological zonation and the interactions of the pests and their biotic and abiotic environments are considered here, with a focus on hosts and climate so that an assessment is based on availability of both host material and suitable climatic conditions.*

*If a pest's distribution is likely to be limited by frosts and/ or low winter temperatures first consider which hardiness zones the pest currently occurs in outside of the risk assessment area. Next consider the area occupied by hosts in relevant hardiness zones within the risk assessment area.*

*If a pest's distribution is likely to be limited by a lack of accumulated temperature, e.g. low summer temperatures, first consider where the pest occurs in terms of accumulated degree day zones outside of the risk assessment area. Next consider the area occupied by hosts in relevant degree day zones within the risk assessment area.*

*Recall that hardiness maps and accumulated degree day maps are based on 10 or 30 year averages. In reality the areas of each zone vary year to year.*

*Taking into account the area of suitable climate and availability of host plants judge what area the pest could potentially establish in.*

*In addition to climate and host, many other factors can be taken into account when assessing likelihood of establishment. ISPM 11 lists other factors to consider, e.g. biotic factors such as the reproductive strategy of the pest, whether the species is polymorphic and the degree to which the pest has demonstrated the ability to adapt to conditions like those in the risk assessment area; the minimum population needed for establishment; competition and natural enemies. Abiotic factors in the environment such as soil type could also be important.*

*Where applicable, practices employed during the cultivation/production of the host crops should be compared to determine if there are differences in such practices between the risk assessment area and the origin of the pest that may influence its ability to establish. Pest control programs already in the risk assessment area which reduce the probability of establishment should be taken into account. Pests for which control is not feasible should be considered more likely to establish than those for which treatment is easily accomplished. The probability of establishment in a protected environment, e.g. in glasshouses, should also be considered.*

*Contracting parties to the IPPC recognise the necessity for preventing the international spread of plant pests and their introduction into endangered areas (FAO, 1997). The IPPC defines "endangered area" as "an area where ecological factors favour the establishment of*

---

<sup>6</sup> Establishment; Perpetuation, for the foreseeable future, of a pest within an area after entry (ISPM No. 5, IPPC, 2007).

a pest whose presence in the area will result in economically important loss". For the purposes of EFSA, assessors are advised to interpret endangered area as "the area where ecological factors favour the establishment of a pest whose presence in the area will result in harmful consequences to cultivated and managed plants<sup>7</sup> and/ or the environment".

If the risk assessment has been initiated by a review of phytosanitary policy where the pest is already present and perhaps widespread in an area, the likelihood of pest establishment should focus on those parts of the area where the pest does not occur.

Fill out table 3.01 by considering how much of the HOST area within the pest risk assessment area is suitable for the pest's establishment taking into account relevant factors such as where host distribution overlaps with suitable climatic conditions (e.g. plant hardiness zones or accumulated degree day zones). For example an assessor could judge that it is 75% likely that 1/3 – 2/3 of the host area is suitable for establishment but 25% likely that between 2/3 and 90% of host area is suitable.

In Table 3.01 distinguish between the host area suitable for establishment and the ENDANGERED AREA which is, for the purposes of EFSA, the area where ecological factors favour the establishment of a pest whose presence in the area will result in harmful consequences to cultivated and managed plants and/ or the environment. In making a judgment regarding the endangered area, the rate of pest population development and any threshold required for harmful pest consequences to materialize within cultivated and managed plants could be taken into account. The magnitude of consequences is considered in Q 3.03 and 3.04.

Sophisticated quantitative environmental modelling could be used to more precisely identify the area where establishment is most likely and to identify the endangered area.

**Information / evidence:** *Provide reasoning then give judgment.*

*Acidovorax citrulli* occurs on all continents although a suggestion that it occurs in Africa (South Africa) has not been confirmed. Whilst the pathogen has been shown to survive on watermelon seeds at 4°C for 26 months (Shirakawa *et al.*, 2003), the disease is generally seen in humid spring or summer months (Bahar & Burdman, 2010). Optimum conditions for disease expression are 55% relative humidity and temperatures between 24°C and 38°C. Hamm *et al.* (1997) considered that the disease may have a significant impact on watermelon production in the southern Columbia Basin, which is dry (<5 cm rainfall), has low relative humidity (46-57%) and high temperatures (29-41°C).

Figure 3.01a shows the world plant hardiness zones (NAPPFASST, 2007) with dotted ellipses showing the approximate occurrence of *Acidovorax citrulli* in relation to hardiness zones, based on the map in Figure 3.01 b showing *A. citrulli* distribution. The pathogen appears to occur in plant hardiness zones 5 or 6 up to plant hardiness zones 10 or 11.

With regard to accumulated temperature *A. citrulli* generally does not appear to occur in areas with less than 1,000 day degrees, above a base of 10°C (Figure 3.01c). However, there is insufficient detailed information for the distribution of *A. citrulli* to accurately determine the accumulated temperature required by *A. citrulli* to express disease symptoms.


---

<sup>7</sup> Managed plants are those plants appreciated / valued/ desired by man, whose growth and spread / distribution are modified by human intervention. It would include plants grown in private gardens.

When outbreaks of bacterial fruit blotch are reported in the literature, there is often mention of hot and humid conditions. Map 3.01d shows the approximate occurrence of the pathogen in relation to Köppen-Geiger climate classification.

Maps in figures 3.01d to 3.01g show the approximate occurrence of *A. citrulli* in relation to global distribution and production of various cucurbit crops (3.01d watermelons; 3.01e cantaloupe and other melons; 3.01f cucumbers and gherkins; 3.01g and pumpkins, squash & gourds).

**Figure 3.01a:** World plant hardiness zones




Source: NAPPFAST (2007) [http://www.nappfast.org/Plant\\_hardiness/ph\\_index.htm](http://www.nappfast.org/Plant_hardiness/ph_index.htm)


Dotted ellipses show approximate occurrence of *Acidovorax citrulli* in relation to hardiness zones (based on map 3.01b, below).

**Figure 3.01b:** Distribution of *Acidovorax citrulli*


Source: EPPO PQR (2011)



**Figure 3.01c:** Approximate occurrence of *Acidovorax citrulli* in relation to global accumulated temperature above a base of 10°C




**Figure 3.01d:** Approximate occurrence of *Acidovorax citrulli* in relation to Köppen-Geiger climate classification




Source: Base map at [http://koeppen-geiger.vu-wien.ac.at/pdf/kottke\\_et\\_al\\_2006\\_A4.pdf](http://koeppen-geiger.vu-wien.ac.at/pdf/kottke_et_al_2006_A4.pdf)

It is noteworthy that bacterial fruit blotch occurs in warm temperate and warm tropical humid zones with either hot or warm summers.


**Figure 3.01e:** Global distribution of watermelon production with approximate occurrence of *A. citrulli*




**Figure 3.01f:** Global distribution of production of cantaloupes and other melons with approximate occurrence of *A. citrulli*



**Figure 3.01g:** Global distribution of production of cucumbers and gherkins with approximate occurrence of *A. citrulli*




**Figure 3.01h:** Global distribution of production of pumpkins, squash and gourds with approximate occurrence of *A. citrulli*



To estimate the area of potential establishment in Europe, the area where potential hosts are grown is considered. Maps 3.01 i to 3.01 l show EU production of cucurbits.

**Figure 3.01 i:** EU area of watermelon production



<b>Table 3.01 i: EU area of harvested watermelon 2007 – 2009 (ha)</b>					
EU MS	2007	2008	2009	3 year mean	% of EU area
Romania	25,224	25,930	28,491	26,548	31.1
Greece	17,108	17,000	17,955	17,354	20.4
Spain	16,861	15,674	17,600	16,712	19.6
Italy	11,141	11,091	11,000	11,077	13.0
Hungary	7,900	7,870	6,918	7,563	8.9
Bulgaria	4,572	4,749	5,593	4,971	5.8
Slovakia	393	334	339	355	0.4
Portugal	350	338	370	353	0.4
France	182	186	200	189	0.2
Malta	125	117	124	122	0.1
Austria	10	13	8	10	0.0
<b>sum</b>	<b>83,866</b>	<b>83,302</b>	<b>88,598</b>	<b>85,255</b>	<b>100.0</b>

Source: FAO Stat Crop production (Area harvested)

Visual estimate of the EU27 watermelon area where day degrees exceed 1000

Figure 3.01 j: EU area of cantaloupe and other melon production





Table 3.0 j: EU area of harvested canteloupe and other melon 2007 – 2009 (ha)					
EU MS	2007	2008	2009	3 year mean	% of EU area
Spain	38,688	33,388	31,400	34,492	37.5
Italy	25,947	28,199	22,300	25,482	27.7
France	14,948	15,603	15,504	15,352	16.7
Greece	8,359	8,400	7,296	8,018	8.7
Romania	3,430	3,672	4,328	3,810	4.1
Portugal	3,100	3,150	3,699	3,316	3.6
Hungary	1,100	878	955	978	1.1
Malta	232	224	195	217	0.2
Slovakia	146	126	106	126	0.1
Netherlands	70	73	67	70	0.1
<b>sum</b>	96,020	93,713	85,850	91,861	100.0

Source: FAO Stat Crop production (Area harvested)


**Figure 3.01 k:** EU area of pumpkin, squash and gourd production



EU MS	2007	2008	2009	3 year mean	% of EU area
Italy	17,023	16,582	13,500	15,702	34.0
Spain	8,500	8,000	5,500	7,333	15.9
Romania	5,603	5,278	5,462	5,448	11.8
France	4,714	5,100	5,054	4,956	10.7
Greece	4,804	3,900	3,211	3,972	8.6
Germany	2,238	2,671	2,728	2,546	5.5
Slovakia	1,831	1,744	1,949	1,841	4.0
Poland	1,528	1,547	1,883	1,653	3.6
Hungary	500	530	1,413	814	1.8
Portugal	700	710	730	713	1.5
Austria	365	367	391	374	0.8
Bulgaria	272	260	226	253	0.5
Latvia	400	74	128	201	0.4
Netherlands	200	200	200	200	0.4
Malta	70	66	54	63	0.1
Finland	46	43	46	45	0.1
Denmark	20	19	18	19	0.0
Lithuania	20	13	16	16	0.0
<b>sum</b>	<b>48,834</b>	<b>47,104</b>	<b>42,509</b>	<b>46,149</b>	<b>100.0</b>

Source: FAO Stat Crop production (Area harvested)

**Figure 3.01 I: EU area of cucumbers and gherkin production**



95% quantile of world presence area = 0.302 ha /km<sup>2</sup>

<b>Table 3.01 I: EU area of harvested cucumbers &amp; gherkins 2007 – 2009 (ha)</b>					
EU MS	2007	2008	2009	3 year mean	% of EU area
Poland	21,036	19,960	20,144	20,380	34.6
Romania	12,590	12,986	13,122	12,899	21.9
Spain	7,466	8,286	8,500	8,084	13.7
Germany	3,224	3,086	3,015	3,108	5.3
Greece	2,765	2,100	2,069	2,311	3.9
Italy	2,145	2,065	2,000	2,070	3.5
Slovakia	2,249	2,191	1,198	1,879	3.2
Czech Repub	1,457	1,655	1,641	1,584	2.7
Lithuania	1,187	1,102	1,050	1,113	1.9
Hungary	1,117	978	1,149	1,081	1.8
Bulgaria	850	371	876	699	1.2
France	642	631	617	630	1.1
Netherlands	600	600	600	600	1.0
Austria	405	365	426	399	0.7
Portugal	350	338	397	362	0.6
Latvia	746	166	77	330	0.6
Finland	354	324	288	322	0.5
Sweden	300	300	300	300	0.5
Estonia	283	261	270	271	0.5
Slovenia	119	115	154	129	0.2
Denmark	120	93	120	111	0.2
UK	103	103	103	103	0.2

Table 3.01 I: EU area of harvested cucumbers & gherkins 2007 – 2009 (ha)					
EU MS	2007	2008	2009	3 year mean	% of EU area
Belgium	100	46	43	63	0.1
Malta	26	27	27	27	0.0
Ireland	15	12	11	13	0.0
	60,249	58,161	58,197	58,869	100.0


Source: FAO Stat Crop production (Area harvested)

As noted above there is no very detailed information regarding the environmental conditions favoured by *A. citrulli*. It has been reported that disease outbreaks are favoured by heavy rainfall and windy conditions (which spreads the pathogen) which expresses symptoms during humid warm periods of summer or spring (Wall & Santos, 1988; Schaad *et al.*, 2008).

The temperature requirements for hosts could be considered as proxies for the temperature requirements for the pathogen. Watermelon germinates best between 21°C and 35°C. Ideal growing temperatures for transplants are between 21°C and 27°C during the day and between 16°C and 22°C at night (CABI CPC, datasheet for watermelon).

Combining plant hardiness zone information, accumulated temperature maps, EU cucurbit host maps and information about the (little described) known distribution of the pathogen and disease, the pathogen could probably establish outdoors in southern and central Europe (Figure 3.01 m). Given that there have been outbreaks of bacterial fruit blotch of cucurbits in Greece, Italy and Hungary survival is known to be possible in these areas already. However, whether the weather conditions prior to outbreaks developing were typical for the areas is unknown.

**Figure 3.01 m: EU cucurbit production, plant hardiness zones and accumulated temperature**



Hosts occur outdoors in plant hardiness zones 6 (e.g. Romania, Poland) to 10 (southern Portugal, Spain and Italy). Melons and cucumbers are grown under glass in northern Europe, where it is not appropriate to consider plant hardiness zones.

## Uncertainties regarding environmental suitability


A combination of appropriate temperature and relative humidity are critical in determining whether *A. citrulli* becomes expressed as bacterial fruit blotch of cucurbits. The range of temperature and relative humidity within which disease symptoms appear are not known although optimum conditions have been described as 55% relative humidity between 24°C and 38°C.

### *Comment regarding the endangered area*

The current assessment considers the endangered area as those watermelons growing regions of the EU where accumulated temperature exceeds 1,000 day degrees above a threshold of 10°C (broadly south of the broken line shown in Figure 3.01n).

Detailed modelling using GIS could be performed to identify where exactly in Europe hosts, relative humidity and temperatures combine to favour disease expression.

**Figure 3.01 n:** Köppen-Geiger climate map for Europe.



### *Conclusions*

Large areas of watermelon production in the EU fall within climate zones assumed to be potentially suitable for establishment of *A. citrulli*. However, whether the conditions over the large areas are suitable for disease expression is more uncertain. It is noteworthy that although the disease has been reported from many US States, in most years the disease occurs in relatively few fields, perhaps due to the particular environmental requirements to stimulate outbreaks.

<b>3.01: Environmental suitability</b> (particularly climate and host)				
<b>Rating</b>	<b>Pest is likely to be able to establish in</b> ...	<b>Justification summary</b>	<b>Probability for suitable area <sup>1</sup></b>	<b>Probability for endangered area %</b>
<b>Very low</b>	Less than 10% of host area		-	-
<b>Low</b>	Between 10% and 1/3 of host area		-	-
<b>Medium</b>	Between 1/3 and 2/3 of host area		-	-
<b>High</b>	Between 2/3 and 90% of host area	Some outdoor crops occur in areas where accumulated temp is below 1,000 DD	15 %	15 %
<b>Very high</b>	More than 90% of host area	Outdoor cucurbits in southern EU occur in areas where pathogen could establish	85 %	85 %
	Check sum =		100%	100 %

<sup>1</sup> Spread your judgment according to your belief / evidence

### 3.02 Extent of spread

*Having found a host and established in the PRA area, a pest will need to spread / disperse after introduction. Consider how quickly the pest could spread. For example, take into account its reproductive potential, suitability of the environment and inherent powers of movement. Assessors should take into account the likelihood that spread may not be contiguous and satellite populations may develop at significant distances from the original point of establishment. Such dispersal could occur via biotic or abiotic vectors, wind, water, or, for example, be facilitated via trade or transport links or via others forms of human assistance, such as movement of infected/infested plant material for propagation purposes (seedlings, scions, budwood).*

*Taking into account the time horizon considered within this assessment (see Initiation, 1.5) estimate the area likely to be occupied by the pest at that time<sup>8</sup>. Fill out table 3.02a by estimating the likelihood that the pest would have spread to occupy the given proportion of the host area suitable for establishment within the time period / time horizon considered by this assessment. Then fill out table 3.02b by estimating the likelihood that the pest would have spread to occupy the given proportion of endangered area within the time period / time horizon considered by this assessment.*

*The tables are used to consider the extent of spread in relation to the area where establishment is suitable. Hence it is recognised that fast moving pests with a large area suitable for establishment may be rated below slower moving pests with much smaller areas suitable for establishment. Quantitative spread modelling could be considered to examine spread more precisely.*

*Specify the time period / time horizon considered by this assessment (refer to 1.5).*

#### **Information / evidence:** *Provide reasoning then give judgment*

Spread of the *A. citrulli* pathogen could occur between commercial crops, from crops to other host plants as well as between other host plants. To estimate how quickly the pest can spread, the following were taken into account: the population dynamics (reproductive potential) of the pest, the influence of the environment and the movement of commodities.

Regarding the population dynamics of the pest, Shirakawa & Yokiko (2003) reported that when infested seeds with only 1 cfu of *A. citrulli* were sown and grown under high humidity, there was a high probability of the germinated seedlings being diseased. The percentage of diseased plants was higher under high humidity than low. Besides, when watermelon seedlings were inoculated with  $10^5$  cfu *A. citrulli* /ml (spray inoculation) and then grown at high humidity, the population of the pest on seedlings increased from less than 10 cfu/g fresh weight to  $10^6$ - $10^7$  cfu/g fresh weight, and seedlings were diseased within 2 days of inoculation. When the inoculated seedlings were incubated at low humidity, the rate of multiplication was lower, reaching  $10^4$ - $10^5$  cfu/g fresh weight by 2 or 3 days, and no seedling was diseased (Shirakawa & Yokiko, 2003). Transmission from seed to seedling is high, and concentration of the pest on seedlings is higher than on seeds (Latin & Hopkins, 1995).

---

<sup>8</sup> When assessing the extent of spread, be clear about the scenario being considered e.g. you could be considering a scenario without risk reduction options in place, or a scenario with specific phytosanitary measures that inhibit spread (risk reduction options) in place.

Hamm *et al.* (1997) reported that when symptomatic test plants were transplanted to fields, maturing melons developed bacterial fruit blotch symptoms.

Regarding plant colonization dynamics by *A. citrulli*, studies on melon plants showed that the bacterium colonized different parts of the melon plant over time, depending on its initial location: leaves or seed (Alves *et al.*, 2010).

The rate of spread depends on inoculum level, aggressiveness of the strains, environmental conditions and presence of host plants.

Hopkins (1993) evaluated the rate of spread from a point-source of inoculum in a watermelon field and found that one infected seedling planted during spring in the centre of a 15m x 15m plot (50 plants) resulted in 80% of the plants showing foliar symptoms before harvest, and 45% of fruit infected. The same severity of foliar symptom incidence occurred in another 7.5m x 30m experimental field plot when three plants were initially artificially inoculated. In the latter case, symptoms developed earlier possibly due to higher starting inoculum level (three infected plants instead of one), but fruit infection reached only 5%, possibly due to dry weather conditions. Regarding inoculum level, Rane & Latin (1992) reported that the bacterial fruit blotch epidemic in Indiana in 1989 was most likely introduced through contaminated seed, and the initial level of contamination was estimated to be 1 infected seed in 9,000.

Regarding aggressiveness of the strains, as described in the relevant datasheet (Table 1) one group (Group I) of strains is considered moderately to highly aggressive to a wide range of cucurbit hosts, while another group (Group II) highly aggressive on watermelon and weakly aggressive on other cucurbits (Walcott *et al.*, 2004; Burdman *et al.*, 2005).

Once infection is established, the pest spreads in intercellular spaces of the seedling and initiate water-soaked lesions. In the transplant facilities, the seedlings are raised under warm and humid conditions, usually in a soilless potting medium. These environmental conditions and the presence of susceptible plant tissue in high density (high plant population) favour disease and the rate of its spread can be especially high (Latin & Hopkins, 1995).

Splashing water from rain or overhead irrigation spread the bacteria from infected seedlings (the source of secondary inoculum) to neighbouring seedlings (Hopkins, 1994; Latin & Hopkins, 1995; Latin *et al.*, 1995). More specifically, in glasshouses, the rate of spread of the disease has been linked to the type of irrigation applied (Hopkins, 1994; Latin *et al.*, 1995). Overhead irrigation resulted in higher disease incidence compared to watering plants from the bottom. Besides, in the first case, symptoms occurred in plants located several cm from another diseased seedling, while in the second case symptoms only spread to plants adjacent to the inoculum source (Hopkins, 1994; Latin *et al.*, 1995). However, with the overhead irrigation, daily low humidity was suggested as a critical limiting factor for the spread of the disease (Hopkins, 1994). Subirrigation has been suggested as a viable option for managing splash-dispersal (Latin *et al.*, 1995).

Reduced relative humidities after overhead irrigation diminished disease spread (Latin *et al.*, 1995). However, successful infection can take place with leaf wetness periods of only 30 min at temperatures around 26°C (Latin, 1996).

The rate of disease spread depends also on the developmental stage of the plants. Young fruits, 2 to 3 weeks after flowering, are most susceptible to infection (possibly through splashing water during rain or irrigation) through stomata and blotch symptoms appear

rapidly on fruit 3-7 days later and generally shortly before they ripen (Frankle *et al.*, 1993). After this period, stomata become blocked by the deposition of waxes on the fruit surface and spread is prevented (Frankle *et al.*, 1993; Walcott, 2005). The surface lesions on mature fruits are an additional source of secondary inoculum in the field and they expand rapidly. Fruit may eventually rot in the field and their seeds, infested by contact with contaminated tissues, slip to the soil (Latin & Hopkins, 1995; Bahar & Burdman, 2010).

The role of wind-driven rain and mechanical means in the local spread of the pest has been acknowledged (Hopkins *et al.*, 2000), but no studies have been made to estimate distances of dispersal by wind-driven rain, as it has been done with other pathogens. For instance: a) *Xanthomonas citri* is readily spread by wind and rain within trees or to neighboring trees (up to 32 meters from infected trees in Argentina), but there is also evidence for much longer dispersals cases associated with meteorological events, such as severe tropical storms, hurricanes and tornadoes, where spread reached up to 56 kms in Florida (Stall *et al.*, 1980; Pruvost *et al.*, 1999; Gottwald *et al.*, 2001; Pruvost *et al.*, 2002; Bock *et al.*, 2010), b) *Pseudomonas syringae* pv. *actinidiae* has shown a spread capacity of around 10 km from the initial infected orchards in Italy (Vanneste *et al.* 2011 in press). The only relevant information found in the literature for *A. citrulli* is that a) honeydew fruits (*Cucumis melo* var. *inodorus*) infected with *A. citrulli* were collected in a field in Texas at 0.8 km from a watermelon field where the disease had been reported 9 weeks earlier (Isakeit *et al.*, 1997), b) citronmelon fruits (*Citrullus lanatus* var. *citroides*) infected with *A. citrulli* were collected in a cowpea field in Texas that was nearby watermelon fields where the disease was present, with the closest field at 50 m from the cowpea field (Isakeit *et al.*, 1998).

No vector has been implicated in the dispersal of the pest (besides humans) (Latin & Hopkins, 1995; Bahar & Burdman, 2010). However, the role of honeybees in watermelon seed infestation through blossom inoculation has been suggested (Fessehaie *et al.*, 2005) but further research is needed to clarify this issue.

It is known that the bacterium can overwinter in volunteer seedlings from seeds from infected fruits, infested rind or other infested crop residues as well as cucurbit weed hosts (especially wild citron), all of them constituting a local source of inoculum that may contribute to outbreaks in subsequent watermelon crops (Latin & Hopkins, 1995). However, according to the NPPOs from 13 EPPO countries (NPPO questionnaires), the citron weed is not known to be present in their countries.

Latin *et al.* (1995) studied the survival of the bacterium on plastic transplants production trays and reported that the bacterium survived 63 days on trays containing potting substrate and root debris. Its longevity decreased with increasing storage temperature and it was not detected after treatment of the trays for 5 min with sodium hypochlorite (10% household bleach).

Spread of the pest can also occur through uncontrolled movement of plant propagation material, e.g. through exchanges of seeds and seedlings between homeowners, especially as latent infections are possible. This is generally a common practice. Although such propagation material can constitute a primary source of inoculum, the volume of this material is very to be likely small. However, if commercial crops are in proximity to such sites (*i.e.* there is host continuity), the risk of spreading the disease is likely. This has been observed with other bacterial pathogens of vegetable crops, e.g. *Xanthomonas citri* spreading from backyard trees to commercial orchards (Gottwald *et al.*, 2001).

As for other bacterial pathogens, the use of non-disinfected tools and equipment, and generally failure to keep sanitized measures in field crops or transplant facilities result in spreading the pest.

In addition, as for all bacterial plant diseases, no curative management measures exists, and it is difficult to contain the pest, especially as even few infected seeds in a lot could initiate a disease outbreak and latent infections may escape inspections.

From the maps (Figure 3.01) showing distribution of major commercial crops in EU that can host the pathogen and the maps on hardiness zones and accumulated temperature, it can be concluded that hosts occur outdoor in plant hardiness zones 6 (Romania, Poland) to 10 (Southern Portugal, Spain, Italy). It has been estimated that the possible establishment area is where accumulated temperatures are above 1,000 degree days. (Presence of host plants grown under protection (glasshouses) have not been related to the hardiness zones)

*Experience from outbreaks in USA:* the pathogen was first observed in commercial watermelon production areas in 1989 and since then it has been reported in many states, for examp, Iowa (1989), Delaware (1989), Maryland (1989), Indiana (1990), Oklahoma (1992), Texas (1994), Oregon (1997), Georgia (1999), Illinois (2002). Outbreaks reported in Maryland and Iowa in 1989 have not recurred, suggesting the introduction of the pathogen with infested or infected propagation material is essential for outbreaks occurrence (Latin & Hopkins, 1995). Thus, the magnitude of threat posed by local inoculum is probably much lower than that of infested or infected propagation material (Latin & Hopkins, 1995).

*Experience from Greece:* The pathogen was detected in 2005 and 2006 on watermelon fruits from fields in very distant areas (Macedonia in Northern Greece and Voiotia in Central Greece, respectively), as well as in 2008 on young watermelon plants from a nursery in the area of Ilia in Southern Greece (Holeva *et al.*, 2010). No further detection of the pathogen has been reported ever since in these areas or any other place (Holeva personal communication).

*Experience from Italy:* The pathogen has been detected in Emilia Romagna (Northern Italy) in 2009 and in Sardinia (Southern Italy) in 2011, regions of Italy separated by approximately 500km and the Tyrrhenian Sea (that part of the Mediterranean sea between mainland Italy, Sardinia and Sicily). These outbreaks could have resulted from separate introduction events rather than the result of *A. citrulli* spread from Emilia Romagna to Sardinia.

#### **Uncertainties regarding extent of spread**

- 1) spread capacity studies are not available,
- 2) the role of pollinators in watermelon seed infestation through blossom inoculation has not been clarified,
- 3) systematic monitoring (surveys) of reported disease outbreaks has not been always available as the pathogen is not currently regulated.
- 4) No detailed data are available for the presence of wild host plants in the risk assessment area.

#### **Conclusions**

Based on the biology of the pest and the epidemiological data mentioned above, when no measures are in place, it is evident that the pathogen has the potential to spread rapidly in the possible establishment area. The current distribution of the pathogen (see datasheet)

reveals that that the progress of the spread so far is in two directions: broadening of the host range and global expansion.

In a scenario of no specific phytosanitary measures being put in place and the likely widespread trade of cucurbit propagation material within the EU, it is possible that the pathogen could be spread across large areas where it could establish, over the next five years. Such rapid spread is made possible via trade networks and “super connected” traders within a network. However, it is unknown how likely a super-connected member of a trade network would be to deal in infested seed or seedlings.

Reports from the USA indicate that when outbreaks occur, the disease incidence is often high, e.g. total losses of fruit at harvest, while fields near severely affected crops can suffer 5 to 50 % loss (Latin & Hopkins, 1995). Thus, the endangered area is the same as the area of potential establishment.

<b>3.02a: Extent of spread in area of potential establishment at time horizon</b>			
<b>Rating</b>	<b>Within the time horizon considered the pest is likely to have spread to ...</b>	<b>Justification summary</b>	<b>Probability that given area will be occupied at time horizon <sup>1</sup></b>
<b>Very low</b>	Less than 10% of the area suitable for establishment	Spread in localised area may be fast by natural means (rain splash & wind-blown) but mostly will not spread very far	60%
<b>Low</b>	Between 10% and 1/3 of the area suitable for establishment	Without <i>A. citrulli</i> being listed, the number of incidences in EU already is low and spread has not exceeded 1/3 of host area	40%
<b>Medium</b>	Between 1/3 and 2/3 of the area suitable for establishment		-
<b>High</b>	Between 2/3 and 90% of the area suitable for establishment		-
<b>Very high</b>	More than 90% of the area suitable for establishment		-
	Check sum =		100%

<sup>1</sup> Spread your judgment according to your belief / evidence

<b>3.02b: Extent of spread in endangered area at time horizon</b>			
<b>Rating</b>	<b>Within the time horizon considered the pest is likely to have spread to ...</b>	<b>Justification summary</b>	<b>Probability that given area will be occupied at time horizon <sup>1</sup></b>
<b>Very low</b>	Less than 10% of the endangered area	As 3.02a above	60%
<b>Low</b>	Between 10% and 1/3 of the endangered area	As 3.02a above	40%
<b>Medium</b>	Between 1/3 and 2/3 of the endangered area		-
<b>High</b>	Between 2/3 and 90% of the endangered area		-
<b>Very high</b>	More than 90% of the endangered area		-
	Check sum =		100%

<sup>1</sup> Spread your judgment according to your belief / evidence

## Consequences of pest establishment and spread

### 3.03 Crop consequences (yield and quality)

*Introduced pests are capable of causing a variety of direct and indirect impacts. The remit of EFSA limits assessors to consider the consequences of pest introduction on crop yield and quality (crop consequences / impacts) (3.03) and environmental consequences /impacts (3.04) e.g. impacts on ecosystem services or biodiversity itself. We recognise that other types of impacts, listed in ISPM 11, may also occur.*

*Fill out table 3.03 by taking into account the extent of pest spread within the endangered area up to the time horizon of the assessment, and other factors such as the rate of pest population development and any threshold required for harmful pest consequences to materialize within cultivated and managed plants. Consequences should be estimated taking into account the current situation in the endangered area with respect to the control efforts undertaken by growers /nurserymen/ producers etc. against other pests. Although we recognise that growers may respond by increasing pest management efforts to minimize impacts of a new pest, such additional efforts are not taken into account.*

*If the risk assessment has been initiated by a review of phytosanitary policy where the pest is already present and action is being taken against it, specify whether consequences are being assessed assuming that action is stopped.*

#### **Information / evidence:** *Provide reasoning then give judgment*

Bacterial fruit blotch has become increasingly important in the watermelon and melon industry since 1987 when the disease destroyed entire fields of watermelon crops on Guam and Tinian in the Mariana Islands (Wall & Santos, 1988; Wall *et al.*, 1990). The disease has now rapidly emerged as a serious pathogen of watermelon and melon crops and is a major threat to these industries around the world (Bahar & Burdman, 2010). In warm and wet conditions (optimally 55% relative humidity and temperatures between 24°C and 38°C), the consequences to the most susceptible crops such as watermelons and melons can be severe.

#### Losses in watermelons and melons in the USA

Following the initial reports of losses from whole fields in the Mariana Islands (Wall *et al.*, 1990), losses of up to 50% of marketable fruit have been reported in Florida (Somodi *et al.*, 1991) and other US States (south eastern, mid-Atlantic, and mid-Western states) as the season progressed. More than 500 ha of watermelons in south western Indiana were affected in summer 1989, with nearly 100 ha sustaining losses approaching 90% (Latin & Rane, 1990). In Texas, when there was above-average precipitation in May and early June 1993, watermelon fields in Frio County were affected with approximately 50% of fruit showing symptoms in affected parts of fields (Black *et al.*, 1994). Fields near to severely affected crops can incur 5 to 50% losses according to environmental conditions and the crop growth stage when infection occurs (Latin & Hopkins, 1995).

Disease incidence of 5% on cantaloupe fruit was reported in Georgia (Walcott *et al.*, 2000) and more than 50% of honeydew fruit loss was reported in Texas (Isakeit *et al.*, 1997). In Oklahoma, in 1991 yield losses in marketable watermelon fruit ranged from 10 to 15% in an affected field (Jacobs & Damicone, 1992).

Although the disease has been reported from many US states in most years the disease occurs in relatively few fields, but where it has occurred it has been devastating in many of them. Bacterial fruit blotch was especially widespread in the USA in 1994 when it caused losses across thousands of hectares distributed over at least 10 States.

#### Losses in watermelons and melons elsewhere

When the disease was found in a watermelon production area of 35,000 ha in the Adana Province in the eastern Mediterranean Region of Turkey in 1995, it caused “serious” yield reductions (Nursen, 2008). In 13 commercial watermelon fields surveyed in the area, 30 – 45% of fruit showed symptoms of bacterial fruit blotch (Mirik *et al.*, 2006). In Brazil crop losses on melon were estimated to be 40 – 50%, and up to 100% in some crops.

In China (Xinjiang Province), during an outbreak, the disease incidence on edible seed watermelon fruit was 30% and on seedlings 20% (Ren *et al.*, 2006).

#### Impacts on other hosts

Although primarily regarded as a pathogen of watermelon and melons, *A. citrulli* has been reported causing lesions on pumpkins leading to total collapse of infected fruit in a commercial pumpkin field in Terrell County, Georgia, USA in September 1998 (Langston *et al.*, 1999). In March 1999, in northern Queensland, Australia, *A. citrulli* caused foliar disease symptoms in a commercial crop of cucumber with more than 20% of the crop affected. No symptoms were observed on plant stems or fruits (Martin *et al.*, 1999). In 2008, *A. citrulli* was reported from betel vine in Taiwan where it caused losses of 30-70% (Deng *et al.*, 2010). This is most significant given that betel vine is not a cucurbit.

#### **Uncertainties regarding crop consequences**

Bacterial fruit blotch has been reported from many locations around the world. However, there is no evidence of the disease recurring year after year at the same location. This could be due to variation in use of contaminated seed or seedlings at locations where outbreaks have previously occurred, or due to the lack of the specific (unknown) environmental conditions occurring in the same locations each year. Losses can vary from 5% to 100% depending on when plants or fruit become infected and environmental conditions. Most reports suggest that when conditions are suitable for the disease, consequences are usually “devastating” for watermelon producers.

## Conclusions

<b>3.03: Potential consequences on crops and managed plants</b>			
<b>Rating</b>	<b>Description</b> (if established in the endangered area, the pest ....(descriptions within categories provide guidance, not all descriptions need to be satisfied in each category)	<b>Justification summary</b>	<b>Probability Assignment</b> <sup>1</sup>
<b>Very low</b>	Under existing pest management regimes, the pest is likely to have <b>negligible or no impact</b> on a standing crop and/or stored products.  Yield and/or quality losses would be <b>negligible and within the range of natural variation</b> .		-
<b>Low</b>	Under existing pest management regimes, the pest is likely to have <b>minimal impact</b> on a standing crop and/or stored products.  Yield / quality losses would be <b>minimal</b> .		-
<b>Medium</b>	Under existing pest management regimes, the pest is likely to have a <b>minor to moderate impact</b> on a standing crop and / or stored products.  Yield / quality losses would be <b>moderate</b> .		-
<b>High</b>	Under existing pest management regimes, the pest is likely to have a <b>moderate to severe impact</b> on a standing crop and / or stored products. Thus the pest will not be effectively controlled by actions already applied against other pests by growers.  Yield / quality losses would be <b>moderate to severe</b> .	When disease results from secondary spread, less plants are affected and impacts are reduced.	50 %
<b>Very high</b>	Under existing pest management regimes, the pest is likely to have a <b>severe impact</b> on a standing crop and / or stored products. Thus the pest will not be effectively controlled by actions already applied against other pests by growers.  Yield / quality losses would be severe.	Several reports suggest that when conditions are suitable for the disease, consequences are usually "devastating" for watermelon producers.	50 %
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

### 3.04 Environmental Consequences

The assessment of the potential of a pest to cause environmental damage proceeds by considering the following factors:

- can the introduction of the pest cause permanent (irreversible) significant, direct environmental impacts, e.g. reduced biodiversity, ecological disruption.
- can the pest have direct impacts on endangered/threatened species by infesting/infecting a plant listed in Annex II or IV of the EC Habitats Directive<sup>9</sup> or infesting / infecting a plant which is a key component of a habitat listed in Annex I of the EC Habitats Directive? If the pest attacks other species within the genus or other genera within the family, and preference/no preference tests have not been conducted with the listed plant and the pest, then the plant is assumed to be a host.
- Can the pest have indirect impacts on species that are listed in Annex II or IV of the EC Habitats Directive or on species that are key components of habitats listed in Annex I of the EC Habitats Directive?
- Would the introduction of the pest stimulate chemical or biological control programmes which would disrupt existing biological or integrated systems for control of other pests or have negative effects on the environment e.g. biodiversity (at various levels), reduce population sizes, or increase their fragmentation.

Fill out table 3.04 by considering the likely magnitude of the above impacts, taking into account the extent of pest spread within the endangered area up to the time horizon of the assessment, and other factors such as the rate of pest population development and any threshold required for the pest to cause environmental harm in the environment.

**Information / evidence:** *Provide reasoning then give judgment*

Although there are wild cucurbit hosts, bacterial fruit blotch does not cause any significant harm in the wider environment or cause any impacts that need to be considered here.

#### Uncertainties regarding environmental consequences

None.

---

<sup>9</sup> Council Directive 92/43/EEC (as amended) on the Conservation of natural habitats and of wild fauna and flora. Available at [http://www.central2013.eu/fileadmin/user\\_upload/Downloads/Document\\_Centre/OP\\_Resources/HABITAT\\_DIRECTIVE\\_92-43-EEC.pdf](http://www.central2013.eu/fileadmin/user_upload/Downloads/Document_Centre/OP_Resources/HABITAT_DIRECTIVE_92-43-EEC.pdf)

## Conclusions

<b>3.04: Potential environmental consequences</b>			
<b>Rating</b>	<b>Description</b>	<b>Justification summary</b>	<b>Probability Assignment <sup>1</sup></b>
<b>Very low</b>	None of the above would occur; the pest is only able to establish on crops grown in protected cultivation such as glasshouses or shade houses. Nevertheless, it is assumed that introduction of a non-indigenous pest will have some environmental impact (by definition, introduction of a non-indigenous species affects biodiversity).	No environmental impacts are reported elsewhere.  This is a pest of commercial crops.	100 %
<b>Low</b>	None of the above would occur; nevertheless the pest could establish outdoors and it is assumed that introduction of a non-indigenous pest will have some environmental impact (by definition, introduction of a non-indigenous species affects biodiversity).		-
<b>Medium</b>	One of the above would occur.  <i>However, if effects are relatively small, the potential consequences can be rated Low instead of Medium.</i>		-
<b>High</b>	Two of the above would occur.  <i>However, if effects are relatively small, the potential consequences can be rated Medium</i>		-
<b>Very high</b>	Three or more of the above would occur.  <i>However, if effects are relatively small, the potential consequences can be rated High</i>		-
		Check sum =	100%

<sup>1</sup> Spread your judgment according to your belief / evidence

### 3.05 Potential impact


The potential impact is assessed assuming entry has occurred and takes into account the endangered area occupied at the given time horizon (3.02b) with consequences to crops (3.03) and the environment (3.04) within the endangered area.

Figure 3.05 shows the BBN model output from combining scores of consequences with the area of the endangered area that could be occupied within the time horizon considered by this assessment.

#### Conclusion

The potentially severe consequences (as seen in USA and Turkey) suggest that impact is likely to be “medium” to “very high”. Given the description of damage caused by the pathogen to cucurbit crops, and particularly watermelons, the model output agrees with the judgment of assessors. Nevertheless, further study of how risk elements are combined in the BBN model is justified.

**Figure 3.05:** Graphical representation of potential impact, combining consequences of pest introduction with the area occupied by the pest at the time horizon.



### 3.06 Pest Risk

To assess the risk for a pest that has not yet entered the PRA area, the potential impact (3.05) which assumes entry has occurred, must be combined with likelihood of entry and transfer (2.08).


Figure 3.06 shows the BBN model output from combining scores for likelihood of entry with potential impact to give an overall pest risk profile. The profile shows much uncertainty with some likelihood of risk being minimal (i.e. very low) up to major (very high). Most likely the pest risk is actually medium to high. Taking into account that major impacts can be seen, but are normally only reported in a small area, this seems a reasonable output for the model.

#### Conclusion

*Acidovorax citrulli* is a pathogen of cucurbits, especially damaging to watermelons, which has spread its geographic distribution in recent years and when environmental conditions are suitable, major crop loss can result.

An experimental protocol based on a BBN, which combines elements of pest risk in a systematic way, expresses overall pest risk as a bar chart using the categories “very low” to “very high”. Further research is required to provide robust interpretations for what each of the categories in the ordinal scale mean. At present a mid-range level of risk with wide uncertainty, as shown in Figure 3.06, approximates the opinion of assessors conducting the assessment. Further study of how risk elements are combined in the BBN model is justified.

**Figure 3.06:** Graphical representation of Pest Risk (3.06), combining overall potential for pest entry and transfer (2.08) with potential impact (3.05).



#### 4.0 Uncertainties

Following EFSA Guidance (EFSA, 2010), to ensure transparency in risk assessment, uncertainties should be identified, characterized and documented within all risk assessments. This can show not only which aspects of an assessment are uncertain but the degree of uncertainty and can help identify where further work could usefully reduce uncertainty.

Table 4: Summary of uncertainties identified and further work that could be undertaken to reduce uncertainties		
Section of risk assessment	Uncertainties	Research that would reduce uncertainty
2.0	Pathways	
<b>2.01a</b>	Pest associated	Survey seed production sites for presence of <i>A. citrulli</i> .
2.02a	Survive post harvest	Study practices used and survey for pest.
2.03a	Survive storage	Monitor survival of seed in store
2.04a	Survives measures	Survey quality checks
2.05a	Quantity imported	Collect import data from trade
2.06a	Transfer	Conduct experiments using contaminated seed
<b>2.01b</b>	Pest associated	Survey seedling production sites for presence of <i>A. citrulli</i> .
2.02b	Survive post harvest	Study practices used and survey for pest.
2.03b	Survive storage/transport	Monitor survival of contaminated seedlings during transport
2.04b	Survives measures	Survey quality checks
2.05b	Quantity imported	Collect import data from trade
2.06b	Transfer	Conduct experiments using contaminated seedlings
<b>2.01c</b>	Pest associated	Survey fruit production sites for presence of <i>A. citrulli</i> .
2.02c	Survive post harvest	Study practices used and survey for pest.
2.03c	Survive storage/transport	Monitor survival of contaminated fruit during transport / storage
2.04c	Survives measures	Survey quality checks
2.05c	Quantity imported	Collect more detailed import data from trade
2.06c	Transfer	Conduct experiments using contaminated fruit and waste disposal techniques
3.01	Environmental suitability	Conduct studies in controlled atmosphere chambers to determine the range of conditions required for disease expression.
3.02	Extent of spread	Detailed climate mapping based on findings from 3.01 with models of spread via trade networks to determine extent of spread.
3.03	Crop consequences	Test broad range of EU cucurbit crops, including different varieties.
3.04	Environmental consequences	Survey for wild cucurbit hosts and potential impacts thereon.

#### Conclusion

The bacterial plant pathogen *Acidovorax citrulli* has the characteristics of a quarantine pest for the EU.

## References

(see also Prima phacie datasheet for *Acidovorax citrulli*)

- Alves, A.O., Xavier, A.S., Viana, I.O., Mariano, R.L.R. & Silveira, E.B. (2010) Colonization dynamics of *Acidovorax citrulli* in melon. *Tropical Plant Pathology* 35, 368-372
- Anon (2011) Bacterial Fruit Blotch: A commercial growers guide. American Seed Trade Association. Information sheet. <http://www.amseed.com/pdfs/DiseaseGuide-BFB-English.pdf>11
- Assis SMP, Mariano RLR, Silva-Hanlin DMW, Duarte V (1999) A bacterial fruit blotch caused by *Acidovorax avenae* subsp. *citrulli* in melon in the state of Rio Grande do Norte, Brazil. *Fitopatologia Brasileira* 24(2), p 191.
- Assouline, I., Milshtein, H., Mizrahi, M., Levy, E. & Ben-Ze'ev, I.S. (1997) *Acidovorax avenae* subsp. *citrulli* transmitted by solanaceous seeds. *Phytoparasitica* 25 (2), 117-118.
- Assouline, I. (1996) The watermelon fruit blotch and other diseases caused by *Acidovorax avenae*. *Phytoparasitica* 24, 136–137.
- Babadoost, M. & Pataky, N. (2002) First report of bacterial fruit blotch of watermelon caused by *Acidovorax avenae* subsp. *citrulli* in Illinois. *Plant Disease* 86 (4), 443.
- Bahar, O. & Burdman, S. (2010) Bacterial fruit blotch: A threat to the cucurbit industry. *Israel Journal of Plant Sciences* 58, 19-31.
- Bahar, O., Efrat, M., Hadar, E. Dutta, B., Walcott, R.R. & Burdman, S. (2008) New subspecies-specific polymerase chain reaction-based assay for the detection of *Acidovorax avenae* subsp. *citrulli*. *Plant Pathology* 57, 754-763.
- Bahar, O., Kritzman, G. & Burdman, S. (2009) Bacterial fruit blotch of melon: screens for disease tolerance and role of seed transmission in pathogenicity. *European Journal of Plant Pathology* 123, 71–83.
- Black, M.C., Isakeit, T., Barnes, L.W., Kucharek, T.A., Hoover, R.J. & Hodge, N.C. (1994) First report of bacterial fruitblotch of watermelon in Texas. *Plant Disease* 78(8), 831.
- CABI (2011) *Acidovorax avenae* spp. *citrulli*. Data sheet last updated 08 April 2011. <http://www.cabi.org/cpc/?compid=1&dsid=2676&loadmodule=datasheet&page=868&site=161>
- CABI (2012) *Citrullus lanatus* (watermelon) data sheet. Crop Protection Compendium. <http://www.cabi.org/cpc/?compid=1&dsid=13678&loadmodule=datasheet&page=868&site=161>
- Cai XueQing, Huang YueYing, Yang JianZhen, Chen Jun, Cai GuoLong, Hu FangPing (2005) Pathogen identification of bacterial fruit blotch of watermelon in Fujian. *Journal of Fujian Agriculture and Forestry University (Natural Science Edition)* 34(4), 434-437 (abst.).
- Crall, J.M., Schenck, N.C. (1969). Bacterial fruit rot of watermelon in Florida. *Plant Disease Reporter* 53, 74–75.
- Demir, G. (1996) A new bacterial disease of watermelon in Turkiye: bacterial fruit blotch of watermelon (*Acidovorax avenae* subsp. *citrulli* (Schaad *et al.*, Willems *et al.*). *Journal of Turkish Phytopathology* 25 (1-2), 43-49.
- Deng, W.L., Huang, T.C. & Tsai, Y.C. (2010) First report of *Acidovorax avenae* subsp. *citrulli* as the causal agent of bacterial leaf blight of betelvine in Taiwan. *Plant Disease*, 94 (8), 1065.
- Dutta, B. & Walcott, R. (2010) Comparative evaluation of the survivability of *Acidovorax avenae* subsp. *citrulli* in stored seeds. *Phytopathology* 100, S200.
- EFSA (2010) Guidance on a harmonised framework for pest risk assessment and the identification and evaluation of pest risk management options by EFSA. EFSA Panel on Plant Health (PLH). *EFSA Journal* 8 (2),1495-1561.
- EPPO (2009a). *Acidovorax citrulli*: addition to the EPPO Alert List. *EPPO Reporting Service*, 2009 / 133.
- EPPO (2009b) *Acidovorax citrulli*. [http://www.eppo.org/QUARANTINE/Alert\\_List/bacteria/Acidovorax\\_citrulli.htm](http://www.eppo.org/QUARANTINE/Alert_List/bacteria/Acidovorax_citrulli.htm)
- EPPO (2010) First record of *Acidovorax citrulli* in Italy. *EPPO Reporting Service* 2010/059.
- Evans, T.A. & Mulrooney, R.P. (1991) First report of watermelon fruit blotch in Delaware. *Plant Disease* 75 (10), 1074.
- FAO (1997) International plant protection convention (new revised text approved by the FAO Conference at its 29th Session— November 1997) [https://www.ippc.int/servlet/BinaryDownloaderServlet/13742\\_1997\\_English.pdf?filename=/publications/13742.New\\_Revised\\_Text\\_of\\_the\\_International\\_Plant\\_Protectio.pdf&refID=13742](https://www.ippc.int/servlet/BinaryDownloaderServlet/13742_1997_English.pdf?filename=/publications/13742.New_Revised_Text_of_the_International_Plant_Protectio.pdf&refID=13742).
- Feng, J., Schuenzel, E.L., Li, J. & Schaad, N.W. (2009) Multilocus sequence typing reveals two evolutionary lineages of *Acidovorax avenae* subsp. *citrulli*. *Phytopathology* 99, 913-920.

- Fessehaie, A. & Walcott, R.R. (2005) Biological control to protect watermelon blossoms and seeds from infection by *Acidovorax avenae* subsp. *citrulli*. *Phytopathology* **95** (4), 413-419.
- Frankle, W.G., Hopkins, D.L. & Stall, R.E. (1993) Ingress of the watermelon fruit blotch bacterium into fruit. *Plant Disease*, **77**, 1090-1092.
- Hamm, P.B., Spink, D.S., Clough, G.H. & Mohan, K.S. (1997) First report of bacterial fruit blotch of watermelon in Oregon. *Plant Disease* **81** (1), 113.
- Harighi, B. (2007) Bacterial leaf spot of Christ's thorn, a new disease caused by *Acidovorax avenae* subsp. *citrulli* in Iran. *Journal of Plant Pathology* **89** (2), 283-285.
- Holeva, M.C., Karafila, C.D., Glynos, P.E. & Alivizatos, A.S. (2009) [Acidovorax avenae subsp. citrulli newly reported to cause bacterial fruit blotch of watermelon in Greece](#). *New Disease Reports [http://ndrs.org.uk]* Volume **20**, 13.
- Holeva, M.C., Karafila, C.D., Glynos, P.E. & Alivizatos, A.S. (2010) [Acidovorax avenae subsp. citrulli newly reported to cause bacterial fruit blotch of watermelon in Greece](#). *Plant Pathology*, **59** (4), 797.
- Hopkins, D.L. (1989) Bacterial fruit blotch of watermelon: A new disease in the Eastern USA. *Proc. Cucurbitaceae* **89**, 74-75.
- Hopkins, D.L. (1993) Field spread of bacterial fruit blotch of watermelon. *Phytopathology* **83**, 466.
- Hopkins, D.L., Cucuzza, J.D. & Watterson, J.C. (1996) Wet seed treatments for the control of bacterial fruit blotch of watermelon. *Plant Disease* **80**, 529-532.
- Hopkins, D.L. & Thompson, C.M. (2002) Evaluation of *Citrullus* sp. germ plasm for resistance to *Acidovorax avenae* subsp. *citrulli*. *Plant Disease* **86**, 61-64.
- Hopkins, D.L., Thompson, C.M., Hilgren, J. & Lovic, B. (2003) Wet seed treatment with peroxyacetic acid for the control of Bacterial fruit blotch and other seedborne diseases of watermelon. *Plant Disease* **87**, 1495-1499.
- IPPC (2004) Pest risk analysis for quarantine pests, including analysis of environmental risks and living modified organisms. International Standards for Phytosanitary Measures No. 11, FAO, Rome, 41pp.
- IPPC (2010) Glossary of phytosanitary terms. International Standards for Phytosanitary Measures No. 5, FAO, Rome, 27pp.
- International Seed Federation ISF (2011) Manual of Seed Health Testing Methods. Available online at: [http://www.worldseed.org/isf/ishi\\_vegetable.html](http://www.worldseed.org/isf/ishi_vegetable.html)
- Isakeit, T., Black, M.C. & Jones, J.B. (1998) Natural infection of citronmelon with *Acidovorax avenae* subsp. *citrulli*. *Plant Disease* **82**, 351.
- Isakeit, T., Black, M.C., Barnes, L.W. & Jones, J.B. (1997) First report of infection of honeydew with *Acidovorax avenae* subsp. *citrulli*. *Plant Disease* **81**, 694.
- Jacobs, J.L., Damicone, J.P. & McCraw, B.D. (1992) First report of bacterial fruit blotch of watermelon in Oklahoma. *Plant Disease* **76** (11), 1185.
- Langston, D. (2011) Bacterial Fruit Blotch Summit. Audio transcript of a radio interview between Dean Kyle (host) and David Langston, (UGA Tifton Vegetable Plant Pathologist), broadcast 11<sup>th</sup> October 2011 on Hometown Country Radio WTIF 107.5. <http://extension.uga.edu/multimedia/tifton-farm-chat/2011/10-11-11-bacterial-fruit-blotch-summit.html>
- Latin, R.X. (1996) Compendium of cucurbit diseases. (Eds. Zitter, T.A., Hopkins, D.L. and Thomas C.E.) APS Press, St. Paul Minnesota, USA.
- Latin, R.X. & Hopkins, D.L. (1995). Bacterial fruit blotch of watermelon. The hypothetical exam question becomes reality. *Plant Disease* **79**, 761-765.
- Latin, R.X. & Rane, K.K. (1990) Bacterial fruit blotch of watermelon in Indiana. *Plant Disease* **74**(4), 331.
- Lessl, J.T., Fessehaie, A. & Walcott, R.R. (2007) Colonization of female watermelon blossoms by *Acidovorax avenae* ssp. *citrulli* and the relationship between blossom inoculum dosage and seed infestation. *Journal of Phytopathology* **155**, 114-121.
- Magarey, R.D., Borchert, D.M. & Schlegel J.W. (2008) Global plant hardiness zones for phytosanitary risk analysis *Scientia Agricola* (Piracicaba, Braz.) **65** (special issue), 54-59
- Martin, H.L. & Horlock, C.M. (2002) First report of *Acidovorax avenae* subsp. *citrulli* as a pathogen of gramma in Australia. *Plant Disease* **86** (12), 1406.
- Martin, H.L., O'Brien, R.G. & Abbott, D.V. (1999) First report of *Acidovorax avenae* subsp. *citrulli* as a pathogen of cucumber. *Plant Disease* **83**, 965.
- Mirik, M., Aysan, Y., & Sahin, F. (2006) Occurrence of bacterial fruit blotch of watermelon caused by *Acidovorax avenae* subsp. *citrulli* in the Eastern Mediterranean region of Turkey. *Plant Disease* **90**, 829.

- Mora-Umana F, Araya CM (2002) Bacterial spot of fruits of melon and watermelon: integrated management of an emergency. *Manejo Integrado de Plagas y Agroecología* **66**, 105-110 (abst)
- Muñoz, M. & Monterroso, D. (2002) Identification of *Acidovorax avenae citrulli* in watermelon seeds in Nicaragua. *Manejo Integrado de Plagas y Agroecología* **66**, 101-104.
- Nascimeto, A.R.P., Mariano, R.L.R. & Silva, E.I. (2004) [Alternative hosts of *Acidovorax avenae* subsp. *citrulli*]. *Horticultura Brasileira* **33** (3), 511-515.
- Nursen, Ü (2008) PRA Record (version 3 of the Decision support scheme for PRA for quarantine pests) *Acidovorax avenae* subsp. *citrulli*, EPPO doc 09-15575. Turkish PRA conducted in 2004 with some updates in 2008. Available from EPPO website [www.eppo.org/.../09-15575%20TR%20PRA%20Acidovorax%20avenae%20pv%20citrulli.doc](http://www.eppo.org/.../09-15575%20TR%20PRA%20Acidovorax%20avenae%20pv%20citrulli.doc)
- Palkovics, L., Petrőczy, M., Kertész, B., Németh, J., Bársony, Cs., Mike, Zs. & Hevesi, M. (2008) First Report of Bacterial Fruit Blotch of Watermelon Caused by *Acidovorax avenae* subsp. *citrulli* in Hungary. *Plant Disease* **92**, 834.
- Rane, K.K. & Latin, R.X. (1992) Bacterial fruit blotch of watermelon: Association of the pathogen with seed. *Plant Disease* **76**, 509-512.
- Ren, Y.Z., Li, H., Li, G.Y., Wang, Q.Y. & Li, J.Q. (2006) First Report of *Acidovorax avenae* subsp. *citrulli* Infecting Edible Seed Watermelon (*Citrullus lanatus* var. *lanatus*) in China. *Plant Disease* **90**, 1112.
- Sato, M (2009) Bacterial fruit blotch of watermelon in Japan and its seed health testing methods. Proceedings *International symposium on the watermelon, Industrial development and utilization*, p10-16, 2009 [book.tndais.gov.tw/Other/.../2009西瓜節研討會專輯10-16.pdf](http://book.tndais.gov.tw/Other/.../2009西瓜節研討會專輯10-16.pdf)
- Schaad NW, Postnikova E, Randhawa P, 2003. Emergence of *Acidovorax avenae* subsp. *citrulli* as a crop-threatening disease of watermelon and melon. In: Iacobellis NS, Collmer A, Hutcheson SW et al., eds. *Pseudomonas syringae* Pathovars and Related Pathogens. *Maratea, Italy: Kluwer Academic Publishers*, 573–81.
- [Schaad, N.W.](#), [Postnikova, E.](#), [Sechler, A.](#), [Clafflin, L.E.](#), [Vidaver, A.K.](#), [Jones, J.B.](#), [Agarkova, I.](#), [Ignatov, A.](#), [Dickstein, E.](#) & [Ramundo, B.A.](#) (2008) Reclassification of subspecies of *Acidovorax avenae* as *A. avenae* (Manns 1905) emend., *A. cattleyae* (Pavarino, 1911) comb. nov., *A. citrulli* Schaad et al., 1978) comb. nov., and proposal of *A. oryzae* sp. nov *Systematic and Applied Microbiology* **31**, 434-446.
- Schaad N.W., Postnikova E., Sechler A., Clafflin L.E., Vidaver A.K., Jones J.B., Agarkova I., Ignatov A., Dickstein E. & Ramundo B.A. (2009). List of new names and new combinations previously effectively but not validly, published. *International Journal of Systematic and Evolutionary Microbiology* **59**, 923-925.
- Schaad, N.W., Sowell, G. Jr, Goth, R.W., Colwell, R.R. & Webb, R.E. (1978) *Pseudomonas pseudoalcalignes* subsp. *citrulli* subsp. nov. *International Journal of Systematic Bacteriology* **28** (1), 117-125.
- Shirakawa, T., Kikuchi, S, Kato, T., Abiko, K. & Kawai, A. (2000) Occurrence of watermelon bacterial fruit blotch in Japan. *Japanese Journal of Phytopathology*, **66**, 223-231.
- Shirakawa, T., Komiya, Y. & Abiko, K. (2003) Population dynamics of *Acidovorax avenae* subsp. *citrulli* on infested seeds and on subsequent seedlings of watermelon. *Japanese Journal of Phytopathology* **69**, 102-106.
- Somodi, G.C., Jones, J.B., Hopkins, D.L., Stall, R.E., Kucharek, T.A., Hodge, N.C. & Watterson, J.C. (1991) Occurrence of a bacterial watermelon fruit blotch in Florida. *Plant Disease* **75** (10), 1053-1056.
- Walcott, R.R., Gitaitis, R.D. & Castro, A.C. (2003) Role of blossoms in watermelon seed infestation by *Acidovorax avenae* subsp. *citrulli*. *Phytopathology* **93**, 528-534.
- Walcott, R.R. (2005) Bacterial fruit blotch of cucurbits. *The Plant Health Instructor*. DOI: 10.1094/PHI-I-2005-1025-02.
- Walcott, R.R., Castro, A.C., Fessehaie, A. & Ling, K. (2006) Progress towards a commercial PCR-based assay for *Acidovorax avenae* subsp. *citrulli*. *Seed Science and Technology* **34**, 101-116.
- Wall, G.C. & Santos, V.M. (1988) A new bacterial disease of watermelon in the Mariana Islands. *Phytopathology* **78**, 1605.
- Wall, G.C., Santos, V.M., Cruz, F.J., Nelson, D.A. & Cabrera, I. (1990) Outbreak of watermelon fruit blotch in the Mariana Islands. *Plant Disease* **74** (1), 80.
- Walcott, R.R., Gitaitis, R.D. & Castro, A.C. (2003) Role of blossoms in watermelon seed infestation by *Acidovorax avenae* subsp. *citrulli*. *Phytopathology* **93**, 528-534.
- Webb, R.E. & Goth, R.W. (1965) A seed-borne bacterium isolated from watermelon. *Plant Disease Reporter* **49**, 818-821.

Zhao TingChang, Sun FuZai, Wang BingWan, Hui WenGuang (2001) Pathogen identification of Hami melon bacterial fruit blotch. *Acta Phytopathologica Sinica* **31**(4), 357-364 (abst).

Zhao, T., Feng, J., Sechler, A., Randhawa, P., Li, J. & Schaad, N.W. (2009) An improved assay for detection of *Acidovorax citrulli* in watermelon and melon seed. *Seed Science and Technology*, **37**, 337-349.

### **Supporting documentation**

Annex 1: Template questionnaire distributed to EFSA Plant Health Network

## Annex 1: Template questionnaire distributed to EFSA Plant Health Network

Request for Information to support EFSA project CFP/EFSA/PLH/2009/01

### *Acidovorax avenae* subsp. *citrulli*

**EU Member State:** Please indicate which EU Member State you are providing information from

Belgium	<input type="checkbox"/>	Greece	<input type="checkbox"/>	Luxembourg	<input type="checkbox"/>	Romania	<input type="checkbox"/>
Bulgaria	<input type="checkbox"/>	Spain	<input type="checkbox"/>	Hungary	<input type="checkbox"/>	Slovenia	<input type="checkbox"/>
Czech Republic	<input type="checkbox"/>	France	<input type="checkbox"/>	Malta	<input type="checkbox"/>	Slovakia	<input type="checkbox"/>
Denmark	<input type="checkbox"/>	Italy	<input type="checkbox"/>	Netherlands	<input type="checkbox"/>	Finland	<input type="checkbox"/>
Germany	<input type="checkbox"/>	Cyprus	<input type="checkbox"/>	Austria	<input type="checkbox"/>	Sweden	<input type="checkbox"/>
Estonia	<input type="checkbox"/>	Latvia	<input type="checkbox"/>	Poland	<input type="checkbox"/>	United Kingdom	<input type="checkbox"/>
Ireland	<input type="checkbox"/>	Lithuania	<input type="checkbox"/>	Portugal	<input type="checkbox"/>		

Please give the total number of regions/departments/provinces/federal states in your country: 1 Department on Plant Health  
(For example: 12 provinces in The Netherlands, 9 federal states in Austria)

### Surveys

**a)** Do you perform surveys to confirm the pest status (presence or absence) of *Acidovorax avenae* subsp. *citrulli*?

No

Region(s) where surveys are performed	Type of places included in the survey <sup>(1)</sup>	Survey frequency <sup>(2)</sup>	Plant species and plant parts included in the survey <sup>(3)</sup>	Type of survey performed <sup>(4)</sup>

<sup>(1)</sup> Indicate the type of places where surveys are performed, by using the following codes: (N) nursery; (F) field crop.

<sup>(2)</sup> Indicate the frequency of surveys for each region, e.g. every year since 2001, 2 times between 2004 and 2010.

<sup>(3)</sup> Indicate the plant species and plant parts surveyed, e.g. watermelon (seeds, seedlings, fruits).

<sup>(4)</sup> Indicate the type of survey performed, by using the following codes: (V) visual inspections; (L) laboratory testing.

### Outbreaks

**b)** Has an outbreak of *Acidovorax avenae* subsp. *citrulli* ever occurred in your country?

YES  NO  NO DATA AVAILABLE

If YES, please complete the table below:

Region(s) / First year of detection <sup>(1)</sup>	Current distribution in the region(s) <sup>(2)</sup>	Type of places where outbreaks occurred <sup>(3)</sup>	Host plants/ plant parts <sup>(4)</sup>	Source of data <sup>(5)</sup>

<sup>(1)</sup> List all regions/departments/provinces/federal states, etc, where the pest was found to occur and the year of first detection.

<sup>(2)</sup> Indicate the current distribution of the pest in each region, using the following codes: (W) widespread; (L) localized; (O) occasionally present; (D) reported in the past, but no longer present; (E) eradicated; (U) under eradication.

<sup>(3)</sup> Indicate for each region if the pest occurred in nurseries or field crops, using the following codes: (N) nursery, (F) field crop.

<sup>(4)</sup> Indicate the plant species and the plant parts found to be infected, e.g. watermelon/seedlings.

<sup>(5)</sup> Indicate the source of data, by using the following code: (S) survey results; (Gs) general surveillance, which includes data from local government agencies, research institutions, universities, scientific societies, producers, consultants, museums and the general public.

## Annex 1 (contd.)

### Request for Information to support EFSA project CFP/EFSA/PLH/2009/01

#### Wild hosts

c) Does the weed Citron (*Citrullus lanatus* var. *citroides*) occur in your country?

YES  NO  NO DATA AVAILABLE

If YES, which is its prevalence in the country?

WIDESPREAD  LOCALIZED  OCCASIONALLY PRESENT  NO DATA AVAILABLE

Has Citron been found in proximity to cultivated cucurbits?

YES  NO  NO DATA AVAILABLE

#### Imports

d) Does your country import cucurbits (propagation plant material and plant products)?

YES  NO

If YES, please complete the following table:

Type of plant material of cucurbits imported	Countries of origin	Number of interceptions <sup>(1)</sup>
Seeds		
Seedlings		
Fruits		
other		

<sup>(1)</sup> Indicate the number of interceptions of imported material, due to *Acidovorax avenae* subsp. *citrulli* infection.

#### Measures

e) Are there any measures applied in your country to prevent introduction and spread of *Acidovorax avenae* subsp. *citrulli*? (e.g. phytosanitary checks, chemical treatment, cultural practices, etc)

.....

.....

.....

.....

.....

#### Contact details

f) Please provide your contact details should we need to contact you again.

Name:	
e-mail:	
Phone:	(international code) , area code, number

Many thanks for completing this questionnaire.  
Your contribution to Prima phacie is greatly appreciated.

Please return this completed form to alan.macleod@fera.gsi.gov.uk by July 5<sup>th</sup> 2010